



HCHS/SOL Manual 15

Endpoint Ascertainment Procedures

July 3, 2014 - Version 1.2

Prepared by the HCHS/SOL Endpoints Subcommittee

Study website - <http://www.csc.unc.edu/hchs/>

Endpoint Ascertainment Procedures TABLE OF CONTENTS

1.0	ENDPOINT ASCERTAINMENT PROCEDURES	1
1.1	OVERVIEW.....	1
1.2	SPECIFIC DISEASE ENDPOINTS.....	1
1.3	ASCERTAINING POTENTIAL EVENTS.....	1
1.4	COLLECTION AND ABSTRACTION OF MEDICAL INFORMATION.....	2
1.5	DEATH INVESTIGATIONS.....	4
1.6	REVIEW AND VALIDATION.....	4
1.7	Clinical Site Procedures for Identifying and Processing Events for Endpoint Classification.....	4
1.8	Medical Records Processing.....	9
2.0	ENDPOINT SURVEILLANCE FOR HOSPITALIZED ACUTE MYOCARDIAL INFARCTION	11
2.1	Introduction.....	11
2.2	Event Identification.....	11
2.3	Screening Codes.....	11
2.4	Diagnostic Criteria.....	11
2.5	Cardiac Symptoms and Signs.....	12
3.0	ENDPOINT SURVEILLANCE FOR HOSPITALIZED HEART FAILURE.....	16
3.1	Introduction.....	16
3.2	Event Identification.....	16
3.3	Diagnostic Criteria for Acute Decompensated Heart Failure.....	16
4.0	ENDPOINT SURVEILLANCE FOR STROKE AND TRANSIENT ISCHEMIC ATTACKS	18
4.1	Event Identification.....	18
4.2	Diagnostic Criteria.....	18
4.3	Definitions of TIA and Stroke.....	18
4.4	Definitions of stroke subtypes.....	19
4.5	Ischemic Stroke Classification.....	20
4.6	Ischemic Stroke Classification.....	22
5.0	ENDPOINT SURVEILLANCE FOR EXACERBATIONS DUE TO CHRONIC OBSTRUCTIVE PULMONARY DISEASE OR ASTHMA	22
5.1	Introduction.....	22
5.2	Event Identification.....	22
5.3	Diagnostic criteria for pulmonary events.....	23
6.0	ENDPOINT SURVEILLANCE FOR PREGNANCY RELATED COMPLICATIONS	27
6.1	Introduction.....	27
6.2	Event Identification.....	27
6.3	Physician review of events.....	28
6.4	Diagnostic criteria for pregnancy related complications.....	28
7.0	ENDPOINT SURVEILLANCE OF FATAL EVENTS.....	31
7.1	Event Identification.....	31
7.2	Diagnostic Criteria.....	31
7.3	Fatal Coronary Heart Disease Events.....	31
7.4	Fatal Pulmonary Events.....	33
8.0	EVENT CLASSIFICATION COMMITTEE.....	33
8.1	Introduction.....	33
8.2	Review Process.....	33
8.3	Disagreement Resolution.....	33
8.4	Confidentiality.....	35

REFERENCES.....36
APPENDIX A. MINNESOTA CODE ECG CRITERIA.....38
APPENDIX B. GLOSSARY OF KEY EVENT DATA COLLECTIONS TERMS41
APPENDIX C. DATA COLLECTION FORMS REQUIRED FOR ASSESSING ENDPOINTS:.....42
APPENDIX D ABSTRACTION FORMS.....61
APPENDIX E REVIEWER FORMS91
APPENDIX F - EVENT SUMMARY FORMS96

1.0 Endpoint Ascertainment Procedures

1.1 Overview. The mission of the Endpoints Ascertainment and Classification Committee is to design and implement a system for event ascertainment, including review and validation of a variety of endpoints to facilitate the classification of incident events occurring among the Hispanic Community Health Study / Study of Latinos (HCHS/SOL) cohort participants. The specific objectives of the endpoints procedure manual are to clearly describe how the HCHS/SOL will:


- identify acute myocardial infarction, stroke, heart failure, asthma and chronic obstructive pulmonary disease (COPD) events that have required hospitalization following the initial examination;
- identify acute exacerbations of asthma or COPD requiring emergency department (ED) care; and
- review and evaluate clinical information collected from medical records from hospitals and emergency departments to classify each event type.

The identification and classification of health events in the HCHS/SOL outlined in this manual follows standard principles of population-based cohort surveillance. These principles include ascertaining potential events, gathering medical information about these events, and reviewing collected data to validate the types of events of interest. The aim of surveillance of the HCHS/SOL cohort is to identify all hospitalizations and emergency department visits for each cohort participant (regardless of reason) and validate the diagnosis of all potential coronary, stroke and pulmonary disease events and key pregnancy related complications (PRC) that occur between baseline exam and the subsequent follow up. We will also investigate deaths to validate cause of death from cardiovascular and pulmonary disease. The general approach to defining endpoints of interest, ascertainment of potential cases, gathering medical information, and review/validation of events is outlined below. Details of each of these steps are provided in the proceeding chapters of this manual.

1.2 Specific Disease Endpoints. The specific cardiovascular and respiratory disease endpoints of primary consideration in HCHS/SOL are: 1) hospitalizations for myocardial infarction, stroke, heart failure, COPD, gestational diabetes mellitus, pre-eclampsia, eclampsia, and asthma, and 2) emergency department visits for COPD and asthma exacerbations. The cause of death of any cohort participant will also be established and is a primary endpoint. Only cardiovascular or pulmonary disease related deaths will be validated.

1.3 Ascertaining Potential Events. Event surveillance of the cohort uses information obtained from the annual phone follow-up interview. When the annual follow-up interview indicates that the participant has either died, been admitted to a hospital, or been seen in an emergency department, mechanisms to obtain the appropriate medical records or death certificate are initiated. Fatal events are also ascertained from review of vital statistics lists and obituaries for the state in which the community is located. The HCHS/SOL records the occurrence of all hospitalizations and all emergency department visits and captures the discharge diagnosis and procedure codes (ICD-9 codes) but only conducts detailed investigations for the selected kinds of medical events noted above. Detailed investigation of recalled hospitalized and emergency department events will be triggered initially by the reported reason for the event but verified by the presence of certain discharge diagnoses or procedure codes (see Figure 1 below). Presence of certain presenting symptoms will trigger investigation of emergency department or emergency medical services (EMS) records. Death investigation will be triggered by certain underlying cause of death codes on the death certificate; and investigation out of hospital deaths will include interviews with next of kin and mailing questionnaires to appropriate physicians, medical examiners or coroners. See Chapter 7 for Endpoint Surveillance of Fatal Events.

Figure 1. Summary of event investigation based on initial reason for hospitalization or emergency department visit as reported by cohort participant



1.4 Collection and Abstraction of Medical Information. A detailed abstraction form specific to the type of event will be used by trained staff at the coordinating center to collect relevant data from medical records of eligible events. Copies of discharge summaries, history and physical, electrocardiograms, echocardiography reports, neuro-imaging reports, consult reports and other pertinent documents will be obtained by field center staff and sent to the coordinating center for abstraction. The type of records from the medical chart to be sought, copied, and sent to the coordinating center is summarized in Table 1. Abstractors follow detailed question by question instructions (QXQ) for the standardized abstraction of medical record information to a database. Abstractors are trained and certified. A brief summary (1-2 pages) of information abstracted from these materials (the Event Summary Form or ESF) will be provided to the Event Classification Committee (ECC) for their review when classifying the event. In addition, copies of selected portions of the materials from the medical record will be provided to the Events Classification Committee members for their use in determining the final event classification of each event.

Table 1. Documents requested for endpoint classification by type of endpoint, determined by responses to self report by cohort participants on annual follow up phone interviews (Annual follow up form item HOE4a)

Medical Record Materials	MI (HOE4a=0)	Chest pain (HOE4a=1)	Heart failure (HOE4a= 2)	Stroke (HOE4a=3)	COPD (HOE4a=6)	Asthma (HOE4a=7)	PRC (HOE4a=9)	Other (HOE4a=8) HOSP only or plus ED
Coding summary with ICD-9-CM codes	√	√	√	√	√	√	√	√
Discharge summary	√	√	√	√	√	√	√	√
History and physical	√	√	√	√	√	√	√	√
Vital signs, weight							√	
12 lead ECG reports (all)	√	√	√	√	√	√		
Cardiac biomarker report	√	√	√		√	√		
Emergency Department report	√	√	√	√	√	√	√	
Cardiac catheterization report	√	√	√					
Arteriogram reports	√	√	√	√				
Stress test report	√	√	√					
Procedure report, delivery or operating room notes	√	√	√	√	√	√	√	
Spirometry report					√	√		
Pulse oximetry report					√	√		
Arterial blood gas report					√	√		
Discharge medication report				√	√	√	√	
Laboratory report	√	√	√	√	√	√	√	
Chest X-ray report			√		√	√		
CT scan report				√	√	√		
Echocardiology report	√	√	√	√	√	√		
RVG or MUGA report	√	√	√		√	√		
Nuclear studies report				√				
MRI report				√				
Lumbar puncture report				√				
Ultrasound report (other than echocardiogram)				√				
Doppler flow study report					√	√		
Carotid studies report				√				
Pulmonary angiography					√	√		
Isotope scan								
Lung scan					√	√		
Autopsy or Medical examiner report	√	√	√	√	√	√	√	

1.5 Death Investigations. A death certificate form will be completed for all eligible fatal events. For in-hospital deaths, data collected on the death certificate form and the hospital abstraction form will be combined for use by the ECC for review and event classification.

Deaths occurring outside the regular acute care hospital are categorized as out of hospital deaths. This includes persons dead on arrival at acute care hospitals, and those dying in outpatient departments or emergency rooms, or admitted without vital signs. For out of hospital deaths meeting underlying cause of death code criteria, information is sought from the decedent's family and physician within 6 months after death. The former is contacted by telephone and the latter by mailed questionnaire. Often the informant is the spouse or other family member of the decedent. Information provided by the informant and the physician is combined for use by the ECC for review and event classification.

1.6 Review and Validation. Diagnostic information obtained through abstraction of the medical record combined with documents copied from the medical records are de-identified and prepared for review by members of the ECC. Cause of death ICD-10 codes obtained through abstraction of the death certificate are combined with supplemental information from informants (in the case of out of hospital deaths) and prepared for review by members of the ECC. ECC members complete an event classification form which indicates their judgment as to the diagnosis of the events. ECC reviewers are trained and certified to follow standardized rules and case laws when determining the final event classification of each case.

1.7 Clinical Site Procedures for Identifying and Processing Events for Endpoint Classification
Annual Follow-Up Form (AFU) (See Manual 16 and study web site for example and QxQ)

During the completion of the annual follow-up interview (contact year 1), field center staff will ask the participant whether they had been admitted to a hospital or seen in an emergency department (ED), at any time since their last HCHS/SOL center visit (HOEA item 3). If no events are reported (HOEA item 1 = 'no' or 'unsure') there will be no events to be investigated. If participants respond 'Yes' to HOEA item 1 then they are asked to identify the type of event in HOEA item 4. HOEA item 4 identifies whether the event was a visit to the ED or an admission to the hospital, or both. This information on event type appears on the Event Tracking Report and can be useful in field center efforts to obtain medical records.

Item 4a on the HOEA form asks: *What was the main reason for going to the (insert emergency room or hospital) that day?* The discharge summary and associated summary of all discharge diagnosis and procedure codes are important parts of the record that will always be obtained for all hospitalized or emergency department events. Events reported during the annual follow up interview that involve an emergency department visit without subsequent hospital admission (a stand-alone ED visit) are selected for further investigation as shown Figure 1 above. Emergency department visits without a subsequent hospitalization are investigated if they are suspected to be for COPD or asthma based on the self-reported reason for the visit (from item 4a of the HOEA form).

Event Identification (ID) Numbers

When an event eligible for investigation is reported during an Annual Follow Up interview and the data is entered into the data management system, a unique event ID number is assigned to each reported event by the endpoints management system. This event ID is derived from the cohort ID number and the reported hospitalization by computer algorithm. If more than one eligible event is reported during an annual follow interview for an individual, then the data management system creates new event IDs for each reported event. For example, for a given cohort ID number there may be several event IDs. Information about the

reported event and the associated event ID is automatically imported into the Event Tracking Table (ETT). An example of the ETT is shown in Table 2.

Table 2. HCHS/SOL Event Tracking Table

Event ID	Cohort name	Date of event	Event type	Medical facility/location	Main reason	Event status tracking	Data status	Comment
X60000130102	Doe, J.	03/04/2008	ED only	Memorial Medical Center	Asthma	Requested	02/01/2009	Pending
X60000130101	Doe, J.	03/01/2008	Hospital	Mercy Medical Center	Heart Failure	Requested	02/01/2009	Release requested

The event ID is composed of the participant’s 8 digit study ID number, followed by 4 digits. The first two digits indicate the year of follow-up (re: 01, 02) the last two digits indicate the sequentially numbered events. (re: 01, 02, 03, 04, etc.)

Examples:

Year one follow-up event labeling for San Diego participant id S8XXXXXX:
 S8XXXXXX0101, S8XXXXXX0102, S8XXXXXX0103, S8XXXXXX0104

Year two follow-up event labeling for San Diego participant id S8XXXXXX:
 S8XXXXXX0201, S8XXXXXX0202, S8XXXXXX0203, S8XXXXXX0204

Event Tracking Table/Status Report (DMS generated)

The Event Tracking Table/Status Report stores key ascertainment data abstracted from the AFU interview. Field site Endpoint Ascertainment staff have password protected access to this report, and have the option of allowing access to the Annual Follow-up Interviewer(s) for ease in confirming changes to the AFU, based on record ascertainment. See the study Data Management system User Guide to create password protected access for selected staff.

The table/report has two drop-down menus in the header section of the screen.

1. Select Event Status: lists all open or closed events
2. Sort By: Sorts by event id, event type or event status

The table/report consists of nine (9) columns. The first 6 columns are auto-populated by DMS

- Column 1: Event ID
- Column 2: Name
- Column 3: Event Date—Item 4b from HOE
- Column 4: Event Type—Item 4 from HOE
- Column 5: Medical Facility/Location—Item 4c from HOE
- Column 6: Main Reason—Item 4a from HOE

Column 7—9 require manual entry by the Endpoint Ascertainment Staff.

- Column 7: Event Status (Click for ETR form)

Event Tracking Report Form (ETR) (See appendix XX for example and QxQ).

This record will be used to track the site efforts on obtaining, processing and shipping the medical records for an event under investigation.

Once an event has been listed on the Event Tracking Status Report in DMS, the staff will create an ETR by clicking on “*No ETR*” under column 7.

In order for the ETR form to auto-save in the system you must completely enter the form. (Event ID and Event Date are pre-populated).

This form should be updated as record acquisition work progresses through “pending records request” to “shipping records to the Coordinating Center”.

Study reports developed will use the ETR to give the field centers and steering committee feedback on the progress of end points investigation activities at each site.

All “closed events” (with terminating/final codes: 4, 5 or 9) will be auto-stored in the Closed Event Table, which is accessed by the “Select Event Status” menu on the Event Tracking Table/Status Report.

Column 8: Status Date--Auto populates once the ETR form is entered.

Column 9: Click for VER Form

Verification Form (VERA) (See appendix C4 for example)

When medical records of interest are received for an event, enter the ICD-9 codes appearing on the discharge summary page into the verification form (VERA) by clicking on the equal sign (=) in the last column for the event.

Once the ICD-9 codes have been entered, the DMS will list the medical record materials expected for this event. For each listing, code whether the document of interest was:

1=received, 2=pending, 3=not available

Successful verification of the discharge codes will produce a face sheet for the materials being transferred to the coordinating center. Table 3 which follows on the next page shows the linkage between the ICD-9 code keyed from the face sheet of the medical record on the VERA form and all of the expected medical record information required to adjudicate the event.

Print the VER form and face sheet to send with the documents. Please note, if the face sheet is not pre-populated with the Subject ID, Date of Event and Event ID—contact the DCC and the staff will assist you.

Table 3. Medical Record Materials Required for HCHS/SOL Endpoint Validation of Hospitalized Events* by Discharge Diagnosis Code

Documents requested for HCHS/SOL Endpoints	Hospital discharge diagnosis codes or procedure codes (ICD-9-CM)						
	MI (402.xx, 410-414, 427, 428, 518.4, 00.50-00.54, 00.61-00.66, 35-39, 88.5, 89.49, 99.10)	Heart failure (398.91, 402.01, 402.11, 402.91, 404.01, 404.03, 404.11, 404.13, 404.91, 404.93, 415.0 425.4, 428, 518.4, 786.0x, 786.2)	Stroke (430-438)	COPD (491.0, 491.2, 491.9, 491.22, 492.0, 492.8, 492.21, 493.0, 493.1, 493.2, 493.9, 496, 786.2, 30.3-30.4, 31.1-31.29, 33.20-33.24, 33.26, 33.27, 33.29, 33.72, 33.99, 89.37, 89.38, 93.90-93.99, 96.01-96.05, 96.7, 96.71-96.72)	Asthma (493, 493.X)	PRC (642.0x 642.1x, 642.2x, 642.3x, 642.4x, 642.5x, 642.6x, 642.7x, 642.8x, 648.0x, 648.8x)	Other discharge or procedure codes
Coding summary with ICD-9-CM codes	√	√	√	√	√	√	√
Discharge summary	√	√	√	√	√	√	√
History and physical	√	√	√	√	√	√	√
Vital signs, weights						√	
12 lead ECG reports (all)	√	√	√	√	√		
Cardiac biomarker report	√	√		√	√		
Emergency Department report	√	√	√	√	√	√	
Cardiac catheterization report	√	√					
Arteriogram reports	√	√	√				
Stress test report	√	√					
Procedure report, delivery or operating room notes	√	√	√	√	√	√	
Spirometry report				√	√		
Pulse oximetry report		√		√	√		
Arterial blood gas report				√	√		
Discharge medication report	√	√	√	√	√	√	
Laboratory report	√	√	√	√	√	√	√
Chest X-ray report		√		√	√		
CT scan report			√	√	√		
Echocardiology report	√	√	√	√	√		
RVG or MUGA report	√	√		√	√		
Nuclear studies report			√				

MRI report			√				
Lumbar puncture report			√				
Ultrasound report (other than echocardiogram)			√				
Doppler flow study report				√	√		
Carotid studies report			√				
Pulmonary angiography				√	√		
Isotope scan							
Lung scan				√	√		
Autopsy or Medical examiner report	√	√	√	√	√	√	

* For emergency department only events selected for investigation, request only the ICD-9-CM coding summary and emergency department report
MI = myocardial infarction
PRC = pregnancy related complications of gestational diabetes, , pre-eclampsia, eclampsia
COPD = chronic obstructive pulmonary disease including emphysema, chronic bronchitis

Updating Events

There may be instances when the participant has reported an incorrect or incomplete event date, which is discovered when the medical records are obtained. The AFU or Data Management (DM) staff can correct the appropriate HOEA by line number for the event. It is very important to select and update the precise HOE record of interest by going to the existing entry for the case, and not creating a new entry (which is default for entering HOE forms).

To do this, right click on the “page” icon just to the left of the participant’s HOE form on the menu on the left side of the DMS screen. Use the “jump to” sub-menu to select the appropriate line.

The program that creates the endpoints workflow table incorporates changes to items on the related HOE form overnight. Corrected information should appear once the CHANGE transaction has cleared the processing system.

Events Not Reported During Annual Follow-up Interviews

Additional events, not self-reported at annual follow-up may be identified when medical records are obtained for a self-reported event. These newly discovered events must be assigned event IDs and must be distinguished from the self-reported events on the HOEA form for the contact year in which it took place.

To correctly capture this information: 1) “jump to” the last HOEA line in the sequence, 2) tab to question 4f, and change that from a “No” to a “Yes” which will bring up a blank HOE form to complete, 3) enter the information for the newly discovered event, 4) Create a note log for HOE item 4b, Date of Event, that says: ***“Detected after AFU interview”***, and 5) Save the changes.

If the newly discovered event is eligible for investigation, it will appear in the work panel as a new entry after the system updates the tracking table overnight.

Numbering Events

If a participant has an ED or hospitalized event where he/she was transferred ***directly*** to another facility, this is considered one (1) event only.

If a participant has discrete admission and discharge dates that are not continuous (with intervening days not in the hospital), these are considered two (2) events.

1.8 Medical Records Processing

Medical Record Release Forms and Cover Sheet (See appendix XX for example)

In order to obtain medical records for a specific event, a current and signed medical record release form will need to be sent to each medical facility identified (as indicated on the Event Tracking Status Report) for each event. Medical record release forms are valid for 90 days from the time the patient signs and dates the form. Keeping in mind that a patient may have more than one event or may have been seen in more than one institution; it may be helpful to have the participant sign several release forms at one time, or to sign one and leave the date blank. Alternatively, some institutions may release medical records with a copy of the participant's signed HIPAA consent form.

The medical records release form must include the specific dates of the event and the name of the healthcare provider requesting the records. A cover sheet with auto-populated demographics and participant's self-reported reason for event from the AFU will accompany each request for medical records.

When records are received, the field center (FC) determines if the records are sufficient to ascertain the event. The first step is to directly compare what was received to what was requested.

ICD Codes (International Classification of Diseases)

Obtaining discharge diagnosis and procedure ICD codes for all events is critical for the standardized ascertainment of potential events. Originally constructed to provide comparable international data on causes of death, today it is used in many countries for coding hospital discharge diagnoses for billing purposes. For both hospitalizations and ED visits, an ICD code is assigned for each diagnosis. Usually there is one *primary* discharge diagnosis/ICD Code for each hospitalization/ED visit, but there may be several *secondary* diagnoses listed as well. The secondary diagnoses may include old and new diagnoses. Usually an ICD summary page is included in medical records for any event and this is often on the face page. If this is not received for a specific event, the FC will need to follow-up with the medical records department to obtain it before the event can be sent to abstraction and processed further. classification . It is necessary to obtain and send the ICD summary page to the Data Coordinating Center (DCC) as part of the medical records package for each identified event. If the ICD-9 summary is not available after exhausting efforts at the medical records department, an attempt to obtain it through the hospital's billing department should then be pursued. If attempts to obtain the ICD-9 summary page are unsuccessful, code the preferred terms used in the discharge summary using the online reference provided below:

ICD9 CODES ONLINE: <http://icd9cm.chrisendres.com/index.php>

If the field center is coding diagnoses without the summary page from medical records, the coder should indicate such on the VER form, Item 2, by entering "3" (unavailable).

De-Identifying and Labeling Medical Records

In order to comply with the de-identification rules for research conducted under HIPAA, field centers are to mask or de-identify the following items on the medical records.

Each page of medical records received must be checked and the following de-identified:

- ✓ Participant name and/or initials
- ✓ Hospital name and street address
- ✓ Institutional letterheads and/or logos
- ✓ Names of everyone (keep degrees or titles)
- ✓ Telephone numbers
- ✓ Medical record numbers
- ✓ Health plan ID numbers
- ✓ Account numbers
- ✓ Social security number
- ✓ Electronic mail addresses
- ✓ Web addresses or URLs, IP addresses

Prior to de-identifying, each page of the medical record must be labeled with the event ID and the date of the event. This may be done by placing a pre-printed label on each page, or writing the information on each page. Blinded materials should be scanned using the document scanner provided for HCHS study use. Scanned medical records should be output from your scanner or workstation to the network server. Use output file type options for a OCR text formatted PDF file that would be named using the event ID of the case (e.g. B12345670201.PDF)

De-identification of hardcopies may be done using a regular point black Sharpie or similar marker. If masking is performed on electronic records then use Adobe Acrobat to redact text of PDFs using a light gray colored overlay (not white, or deep black).

Transfer of Electronic Medical Records to the Data Coordinating Center

Scanned endpoints materials should be uploaded weekly to the HCHS coordinating center. More than one event may be batched in an upload with each of the separate PDF files uniquely named using the event ID of the case. The face sheet from the VER should be appended electronically in front of the first page of the scanned medical records. Weekly secure FTP upload requests for endpoints materials will be made by CC staff. Never email medical records to the CC, always use the secure FTP transfer mode instead. Electronic transfer is always preferred to postal mail delivery.

Shipping Printed Materials to the Data Coordinating Center

If hardcopy of records needs to be transmitted then bundle materials for each participant event separately. The face sheet printed from the VER form must appear as the first page for each event packet. More than one event may be shipped together as long as each event has been accurately labeled. All medical records must be shipped by a service providing a tracking mechanism, such as Federal Express. Medical records may be shipped using 2nd or 3rd day delivery option. Next day delivery is not necessary and is more expensive. Remember to maintain the tracking number of the package(s) until notification of receipt by the Data Coordinating Center has been emailed to you.

Please mail medical record packet(s) to:
 Barbara Brown (phone: 919-962-3092)
 HCHS/SOL Event Receiving
 Collaborative Studies Coordinating Center
 137 East Franklin St., Suite 203
 Chapel Hill, NC 27514

In the event that a record received at the CSCC has not been de-identified, the FC will be notified of this error, the record at the CSCC will be appropriately destroyed and a de-identified copy will need to be sent from the FC.

2.0 Endpoint Surveillance for Hospitalized Acute Myocardial Infarction

2.1 Introduction

The aim of surveillance of the HCHS/SOL cohort in regard to acute myocardial infarction is to identify all hospitalizations for each participant and validate the diagnosis of all potential coronary events. The criteria outlined below were created to be comparable to that used by MESA. Ascertainment and validation of all out-of-hospital fatal events that are potentially cardiac-related are also completed (see Endpoint surveillance of fatal events, Section 7 of this manual).

2.2 Event Identification

If a participant reports any hospitalization, field center staff requests the discharge summary, discharge diagnoses an associated ICD-9-CM codes, and any related test results and progress notes from the hospital (see Table 1). A recent signed consent is required by most hospitals in order to release records (see earlier in this section for more information about consents). Once the record is received, field center staff matches the reported hospitalization to the actual record and, if discrepancies are found, re-contacts the participant to resolve these differences. If the event involved a transfer to another hospital or other health-care facility, field center staff obtain all pertinent records from all institutions. Transfers are considered together as one potential investigation.

Endpoints staff at the Coordinating Center reviews the ICD-9-CM codes and, if necessary, the discharge summary, to complete the Events Eligibility form, which determines whether the hospitalization is eligible for further detailed record abstraction. (Please see Table 3.1 for a list of eligible ICD codes.) Pertinent parts of the hospital medical record will be copied and sent to the coordinating center for central abstraction. Components of the chart are scanned at the coordinating center and stored for use by the Endpoints Review committee.

2.3 Screening Codes

Events with CPT procedure code 35 or ICD-9 discharge diagnosis codes 250, 390–459, 745–747, 794.3, 798-799 are eligible for detailed abstraction by coordinating center staff for myocardial infarction. Events without these target codes, yet upon review the discharge summary by coordinating center abstractors contain evidence of eligible conditions an acute myocardial infarction event are also eligible for detailed abstraction..

2.4 Diagnostic Criteria

2.4.1 Myocardial Infarction. Myocardial infarction is defined as the death of part of the myocardium due to an occlusion of a coronary artery from any cause, including spasm, embolus, thrombus or rupture of a plaque. The algorithm for classifying MI includes history of chest pain,, evidence from cardiac biomarkers, and ECGs, The criteria to be used in HCHS/SOL was designed to be comparable with that used in the MESA Study. Additional event classification elements (e.g. anatomical location of MI) not available in MESA were also incorporated into the HCHS/SOL MI validation process.

The definition includes MI that occurred during surgery/procedure and MI aborted by thrombolytic therapy or procedure. The differentiation of definite vs. probable MI will be made based on the criteria described below. These criteria are summarized in Table 4.

Table 4. HCHS/SOL Diagnostic Criteria for Hospitalized MI, adapted from MESA

ECG Pattern*	Abnormal Enzymes**	Equivocal Enzymes**	Incomplete Enzymes**	Normal Enzymes**
Chest Pain PRESENT:				
Evolution of Major Q-Wave	Definite MI	Definite MI	Definite MI	Definite MI
Evolution of ST <u>Elevation</u> with or without Q-wave <u>Or</u> New LBBB	Definite MI	Probable MI	Probable MI	No MI
Evolution of ST-T Depression/inversion alone <u>Or</u> Evolution of Minor Q-waves alone	Definite MI	Probable MI	No MI	No MI
Single ECG with Major Q-Wave <u>Or</u> Single ECG with LBBB, described as new	Definite MI	Probable MI	No MI	No MI
Normal, Absent, Uncodable, other	Probable MI	No MI	No MI	No MI
Chest Pain ABSENT:				
Evolution of Major Q-Wave	Definite MI	Definite MI	Definite MI	Definite MI
Evolution of ST <u>Elevation</u> with or without Q-wave <u>Or</u> New LBBB	Definite MI	Probable MI	No MI	No MI
Evolution of ST-T Depression/inversion alone <u>Or</u> Evolution of Minor Q-waves alone	Probable MI	No MI	No MI	No MI
Single ECG with Major Q-Wave <u>Or</u> Single ECG with LBBB, described as new	Probable MI	No MI	No MI	No MI
Normal, Absent, Uncodable, other	Probable MI	No MI	No MI	No MI

* ECG categories are listed in Appendix 6. Definite indicates definite MI; Probable, probable MI; and No, no MI. Classification of case is at highest level allowed by combinations of 3 characteristics (cardiac signs and symptoms, ECG findings, biomarkers).

**Abnormal and Diagnostic = Adequate set and $\geq 2 \times \text{ULN}$. Adequate are two sets at least 6 days apart at the time of the event.

Equivocal = Present but not diagnostic.

Incomplete = Not available for the time of the event.

2.5 Cardiac Symptoms and Signs

2.5.1 Chest Pain. Chest pain is defined as an episode of pain, tightness, pressure or discomfort in the chest, arm or jaw. If there is a clear non-cardiac cause, chest pain is considered to be absent. Duration of pain is not a part of the chest pain criterion. Adjudicators will assess for the presence of chest pain documented around the time of the potential MI. For events where the MI occurs after admission, chest pain at the actual time of the acute event being adjudicated is relevant (rather than, for example, chest pain prior to admission).

2.5.2 Cardiac Biomarker Criteria. The 2003 AHA Scientific Statement defined positive biomarkers as “the 99th percentile of the distribution in healthy populations or the lowest level at which a 10% coefficient of variation can be demonstrated for that laboratory.” However, as of fall, 2004, this recommendation has not been consistently implemented by manufacturers and upper limits of normal range from the 95th percentile to the 99th percentile, with coefficients of variation difficult to ascertain at these levels. Many manufacturers continue to include an “indeterminate” range, often from the upper limit of normal to some higher value. This indeterminate range is not recommended to be of interest for

determination of MI in epidemiologic studies according to the 2003 Position Statement, and will not be of interest to HCHS/SOL. Because of the continued inconsistency of reporting of the 99th percentile and the 10% coefficient of variation, HCHS/SOL will follow the practice of the MESA summarized in Table 3.

In the event that the actual laboratory values are not included in the medical record then biomarker results reported in physician notes are acceptable, as long as actual values are reported. Reports of biomarkers being “positive” or “negative” will not be sufficient. Use Table 1 to classify cardiac biomarkers. To summarize Table 1, equivocal biomarkers are between “above normal” and twice the Upper Limit of Normal (ULN), whereas “abnormal” biomarkers are greater than twice the upper limit of normal. When there has been muscle trauma, liver trauma, or hemolysis then positive enzymes are downgraded to equivocal. These criteria apply as long as the patient has not had Coronary Artery Bypass Surgery (CABG) or Percutaneous Transluminal Coronary Angioplasty (PTCA) in the previous 24 hours. In that case, see foot notes to classify.

Table5. Algorithm to classify cardiac enzymes as abnormal, equivocal, or normal

Enzyme Value	If: a) no known muscle trauma or hemolysis, and 2) no PTCA or CABG in previous 48 hours*	If Muscle trauma or liver trauma or hemolysis exists then:
CK-MB = “present”, where only present or absent provided	Abnormal	Equivocal
CK-MB ≥ 2X ULN (upper limit of normal)	Abnormal	Equivocal
CK-MB** ≥ 10% Total CK, if no ULN is given	Abnormal	Equivocal
Total CK ≥ 2X ULN <u>and</u> LDH ≥ 2X ULN	Abnormal	Equivocal
LDH1: LDH2>1	Abnormal	Equivocal
LDH1 ≥ 2X ULN if LDH2 is missing	Abnormal	Equivocal
Total CK ≥ 2X ULN <u>or</u> LDH ≥ 2X ULN	Equivocal	Normal
Normal < Total CK < 2X ULN <u>and</u> Normal < LDH < 2X ULN	Equivocal	Normal
5% Total CK < CK-MB** < 9 % Total CK <u>or</u> CK-MB “weakly present”	Equivocal	Equivocal
Normal < CK-MB < 2X ULN	Equivocal	Equivocal
Normal < LDH1 < 2X ULN	Equivocal	Equivocal
Data present, but insufficient for criteria	Incomplete	Incomplete
Normal< Troponins < 2X ULN	Equivocal	Equivocal
Troponins > 2X ULN	Abnormal	Abnormal
Troponins < ULN	Normal	Normal
CK-MB < ULN	Normal	Normal
All other results	Normal	Normal

*If PTCA then abnormal in first 48 hours if Troponins or LDH1 or CK or CK-MB>3X ULN; equivocal if 1-3X ULN. If CABG then abnormal in first 48 hours if troponins or LDH1 or CK-MB>5X ULN; equivocal if 1-5X ULN.

**CK and CK-MB must be in same units for this criterion

2.5.3 ECG Criteria. The ECG criteria for HCHS/SOL are based on the Minnesota Code system of classification which is outlined in the Minnesota Code ECG Criteria shown in the appendix. This is also the same criteria used in MESA. The reviewers will interpret serial tracings; the following ECG tracings are identified and provided for this purpose: the first, second, third and last ECGs obtained from the hospital admission.

The evolution of ECG findings may be demonstrated (1) between the ECG(s) associated with the event or (2) between a previously recorded ECG and the event ECG(s). In cases in which only a single event ECG is available, an evolving diagnostic ECG pattern cannot be recorded. In order to ascertain MI by ECG, precise guidelines to determine wave duration and voltage will be determined following the Minnesota Code. HCHS/SOL reviewers will consider the copies of actual ECGs submitted in the case packet and will make a clinical reading of the ECG pattern, using the Minnesota Code as a guide. The categories of ECG are: a) Evolution of Major Q-Wave, b) Evolution of ST-T Elevation with or without Q-Wave, c) New LBBB, d) Evolution of ST-Depression/T wave inversion alone, e) Evolution of Minor Q-Wave alone, f) Single ECG with Major Q-wave, g) Single ECG with LBBB, described as new, and h) Absent, Uncodable or Other ECG.

2.5.4 Subclassification of definite or probable MI: How to define MI type, location, and whether related to a procedure

MI type: For events classified as definite or probable MI, describe the type of MI as either transmural or subendocardial or undetermined (unsure). Record TRANSMURAL if there is ST elevation or a resulting Q wave note on the ECG. Record SUBENDOCARDIAL if there is no evidence of ST elevation or a resulting Q wave on the ECGs provided.

MI location: Based on your interpretation of the available ECGs, select the location of MI that best describes this event. General guidelines for defining MI location will be those outlined in AHA/ACCF/HRS recommendations for standardized interpretation of ECG (Wagner et al JACC, 2009;53:1003-1011). This will be discussed and revised further at the reviewer training and this manual modified accordingly. Reviewers are asked to classify the MI location for definite or probable MIs as either: 1) Anterior, 2) Posterior, 3) Inferior, 4) Lateral, 5) Septal, 6) Anteroseptal, 7) Anterolateral, or 9) Unable to determine.

In general, MI locations are defined by the following ECG changes and coronary artery occlusions		
MI location	ECG findings	Usually due to occlusion of:
Inferior MI	ST elevation, Q wave in II, III, and AVF	right or left coronary, depends on which coronary supplies the base of the heart
Lateral MI	ST elevation, Q wave in I and AVL	Circumflex branch of L coronary
Anterior MI	ST elevation, Q wave in V1-V4	Anterior descending branch of L coronary
Posterior MI	“opposite” of anterior MI, ST depression, R wave in V1, V2	R. coronary or one of its branches

Procedure-related MI: Cases classified as either definite or probable MI will be further assessed as to whether or not they are related to a procedure. Cardiac events up to 28 days after a medical procedure or surgery may meet criteria for being procedure-related. If the medical procedure was performed for the treatment of acute ischemia (e.g., angioplasty following the presentation of acute coronary syndrome), that

event should *not* be considered procedure-related (Luepker, et al, 2003). The procedure-related MI category is intended to identify MIs that occurred only after the procedure, and were not already in evolution prior to the procedure. In determining whether the MI was procedure related, answer YES (and choose whether it was a cardiovascular or non-cardiovascular procedure) if you think it is unlikely this MI would have occurred had the procedure not been performed. Answer YES, **CARDIOVASCULAR PROCEDURE** if the MI occurred within 28 days of a cardiovascular procedure or surgery AND in your judgment the MI was a complication (or related to) the procedure. Cardiovascular procedures include: CABG, valve replacement, AICD or pacemaker placement, PTCI, etc. Answer YES, **NON-CARDIOVASCULAR PROCEDURE** if the MI occurred within 28 days of a non-cardiovascular procedure or surgery AND in your judgment the MI was a complication (or related to) the procedure. Non-cardiovascular procedures include all procedures or surgeries that are not cardiovascular. Answer UNKNOWN/UNSURE if you are not certain as to whether the MI was procedure related or not.

2.5.5 Angina For events reviewed that are classified as “no MI” or “unclassifiable, then as a secondary endpoint reviewers are asked to state whether angina was present. Angina is a symptomatic event generally involving ischemic chest, left arm, or jaw pain, though the symptoms may be "atypical." Atypical anginal symptoms can include shortness of breath, exertional dyspnea, epigastric discomfort, and back pain, in addition to pain that is isolated to the arm or the jaw. HCHS/SOL endpoint reviewers categorize angina events as “definite,” “probable,” and “no Angina” based on their clinical judgment in light of the following criteria from the MESA study in answering this question:

- a. Physician diagnosis of angina and receiving medical treatment for angina (e.g., nitrates, beta-blockers, or calcium-channel blockers)
- b. CABG surgery or other revascularization procedure
- c. 70% or greater obstruction of any coronary artery per angiography
- d. Horizontal or down-sloping ST-segment depression or abnormal ST elevation of ≥ 1 mm on exercise or pharmacological stress testing with pain
- e. Scintigraphic or echocardiographic stress test positive for ischemia
- f. Resting ECG shows horizontal or down-sloping ST depression or abnormal ST elevation ≥ 1 mm with pain that is not present on ECG without pain

Given the difficulty in the diagnosis of angina yet the need to standardize its classification as much as possible, HCHS/SOL endpoint reviewers are instructed to follow the guidelines a-d below when recording their answer.

- a. Clear and thorough documentation of symptoms is needed to identify an event as “definite angina.” Even if a test such as an ETT lists “angina” or “chest pain” as its indication, angina should not be ruled as being present unless there is additional, explicit information from the physician regarding symptoms. Likewise, a test showing positive ischemia or the performance of a further procedure (e.g., catheterization) is not enough to rule for angina if other HCHS/SOL criteria are not met.
- b. Only classify an event as angina if it is distinct from an MI.
- c. Reviewers should not classify angina as part of pain symptoms of an MI.
- d. Angina will require clinical symptoms. If there is only a physician diagnosis/treatment, then the diagnosis cannot be ‘definite.’ If there is more than just a physician diagnosis, then the reviewer can assign ‘definite’ instead of ‘probable.’

2.5.6 Revascularization procedure interrupting an MI. Revascularization procedures occurring during the course of hospitalization in general will be documented by the abstractors. However, HCHS/SOL endpoint reviewers are asked to record whether in their judgement an intervention performed early in the clinical presentation of a potential MI may have prevented an MI. In cases where revascularization was performed without clinical symptoms, HCHS/SOL endpoint reviewers will record NO to this item. Reviewers should record YES, if on presentation with chest pain or other MI symptoms, the patient is immediately received a revascularization procedure.

3.0 Endpoint Surveillance for Hospitalized Heart Failure

3.1 Introduction

All cases of hospitalized heart failure among HCHS/SOL participants will be identified through the annual follow up call. All eligible hospitalization will be investigated and processed through the HCHS/SOL Event Classification Committee. Heart failure events resulting in outpatient diagnosis and treatment without hospitalization will not be identified and reviewed by HCHS/SOL event reviewers. See nonfatal outpatient event surveillance for details.

3.2 Event Identification

Events to be investigated for hospitalized heart failure include those with the following target ICD-9 discharge diagnosis codes: 402, 404, 415, 416, 425, 428, 518.4, and 786. Specified components of the medical record from eligible events will be copied and sent to the coordinating center for processing. Data from the medical record of hospitalizations with these discharge diagnosis codes will be abstracted using the HCHS/SOL heart failure abstraction form. Materials from the medical record to be copied and provided to the HCHS/SOL event reviewers include: the first three (3) chest X-ray reports, echocardiography reports, cardiology consult report, discharge summary, and cardiac catheterization report.

3.3 Diagnostic Criteria for Acute Decompensated Heart Failure

The HCHS/SOL criteria for heart failure were adapted from the MESA and the Atherosclerosis Risk in Communities study. HCHS/SOL physician reviewers will determine if the event has a heart failure diagnosis from the provider and whether the patient was treated for heart failure. The reviewer will also determine if there is sufficient evidence to indicate the patient has history of heart failure and whether there was X-ray pulmonary edema or congestion. Evaluation of these items follow MESA guidelines and allow for comparability of heart failure diagnosis between MESA and HCHS/SOL. (See Heart Failure Diagnosis (HFD) Form)

In addition HCHS/SOL physician reviewers categorize acute decompensated HF (ADHF) events as “definite,” “probable,” “no ADHF”, and “unknown” in a manner adapted from the ARIC classification scheme for heart failure.. (See Heart Failure Diagnosis (HFD) Form).

Specifically the HCHS/SOL reviewers are asked to evaluate the evidence for the following items:

- a. Heart failure diagnosed by physician, and treatment provided for heart failure,
- b. Acute decompensated heart failure,
- c. Pulmonary edema/congestion by chest X-ray,
- d. Cardiac imaging study results, (Each of these questions is asked separately and if present then specify, as to whether the finding was by history, or by current imaging.)
 1. Dilated ventricle or
 2. Poor left ventricular function (e.g., low ejection fraction or wall motion abnormalities),

3. Poor right ventricular function, or
4. Left ventricular diastolic dysfunction.
5. If available, the quantitative ejection fraction is specifically provided within a range of choices of ≥ 50 , 40-49, 30-39, 20-29, <20 or unknown.

This approach has the advantage of easily permitting a range of analyses based on definitions of heart failure that include "soft" criteria or various types of "hard" criteria.

In general, the reviewer should examine the original report of a procedure rather than accept references to results of the diagnostic or therapeutic procedures in discharge summaries. If an original full report is not available, a convincing reference to the procedure results in the discharge summary is acceptable.

In addition, HCHS/SOL reviewers evaluate the evidence for or against history of heart failure, severity of heart failure and presence of right-sided heart failure, comorbid conditions, and asymptomatic ventricular dysfunction. These are described in more detail below.

3.3.1 Prior history of heart failure. Reviewers are asked to classify the event to whether participant had a prior history of HF or not. Prior history is relevant as the evaluation of a patient with an established history of HF may be more limited as compared to a patient with a new diagnosis of heart failure. This also allows the HCHS/SOL to classify those with chronic stable heart failure if the answer is YES to prior history of HF and NO for the classification of ADHF.

3.3.2 Severity of the heart failure exacerbation. Reviewers classify the severity of the HF exacerbation for those classified as probable or definite ADHF. Classify the event as SEVERE if treatment with mechanical ventilation, non-rebreather mask, CPAP, hemofiltration, intraortic balloon pump, or thoracentesis was required for management of HF exacerbation. Classify event as MODERATE, if it is clear the event was neither SEVERE nor MILD. Classify event as MILD if this exacerbation could have been managed in the outpatient setting had the patient been an outpatient. In these cases, the primary reason for being hospitalized will likely be something other than HF. If it is unclear as to the severity of the event then classify as UNKNOWN.

3.3.3 Predominantly right-sided heart failure. The most common presentation for ADHF is left sided HF which is dyspnea and pulmonary edema, however symptoms of pure right sided HF do not include dyspnea. In the future when considering criteria for the diagnosis of HF, it may be necessary to differentiate left and right sided HF. Answer YES if the patient had right-sided heart failure symptoms only of lower extremity edema and possible ascites and a normal LV ejection fraction. Answer NO/NR if the patient had both right-sided and left-sided signs/symptoms, or if it is clearly not right sided HF, or it is unclear whether the patient had predominantly right sided failure.

3.3.4 Co-morbid and potentially precipitating factors to ADHF. Acute decompensated HF may be precipitated by multiple conditions. In addition, many of the listed comorbid condition may be caused by the exacerbation. As a result, it can be difficult to tell whether the condition precipitated the event or not. Here we ask reviewers to consider the timeline of occurrence of the condition with the HF exacerbation and based on their judgment state as to whether the condition may have precipitated the HF exacerbation. As stated in the question by question instructions for the HFD form, reviewers indicate all co-morbid conditions that were active during this hospitalization AND may have precipitated the event. The temporal association of between the event and the HF exacerbation should make sense. For example, if the patient came in for surgery and had an exacerbation of HF immediately following surgery due to fluid overload then check YES to fluid overload. If the same patient then had a complicated course and developed a PE many days later then check NO/NR for PE, if it temporal association with the HF

exacerbation is not correct. If a patient presents to the hospital with heart failure AND one of these diagnosis that is active then check YES. For example, if presents with both HF and atrial fibrillation the check YES for atrial fibrillation. Reviewers answer No/Not recorded or YES to all of the following: 1) Myocardial infarction, 2) Atrial fibrillation or atrial flutter, 3) Other arrhythmia, 4) Fluid overload or volume overload- this refers only to either iatrogenic fluid overload OR to excessive drinking of fluids, or renal failure due to missed or inadequate hemodialysis, 5) Medication noncompliance – Medication noncompliance would include refusal of medications for patients in the hospital, but more commonly this would be outpatient noncompliance. Noncompliance includes those patients who did not get their medications due to lack of funds, 6) Pulmonary embolus, 7) Renal insufficiency or failure – use your judgement as to whether renal failure was of a severity that it may have contributed to the HF. This would likely be those who are approaching dialysis and are less responsive to diuretics, 8) Cardiovascular procedure/surgery, 9) Non-cardiovascular procedure/surgery, 10) Pulmonary disease, and 11) Uncontrolled Hypertension – systolic blood pressure >180 at the time of the event.

3.3.5 Asymptomatic left ventricular dysfunction. The focus of the primary outcome in HCHS/SOL is classification of acute decompensated heart failure (adapted from ARIC criteria) and the MESA definition of physician diagnosed and treated heart failure. In addition, for those events that are not classified as “definite” or “probable” ADHF then we ask reviewers to identify those with asymptomatic left ventricular dysfunction, defined here as documented ejection fraction < 50%, but no HF symptoms previously or during this admission.

4.0 Endpoint Surveillance for Stroke and Transient Ischemic Attacks

4.1 Event Identification

A hospitalization is eligible for stroke evaluation and classification by the HCHS/SOL reviewers if it has an ICD-9 procedure code 38-39 or discharge diagnosis codes for cerebrovascular (ICD-9-CM 430-438). If a hospitalization meet procedure or discharge code criteria for stroke or transient ischemic attack (TIA,) the medical records obtained by field center staff and sent to the coordinating center for detailed abstraction using the HCHS/SOL stroke abstraction form. These data will be summarized and made available to the stroke group of the Event Classification Committee. Copies of specific portions of the medical record are also provided to the HCHS/SOL reviewers. The reviewers will follow the criteria below in making a final event classification.

4.2 Diagnostic Criteria

The classification criteria for stroke and TIA for HCHS/SOL were adapted from MESA.

4.3 Definitions of TIA and Stroke

4.3.1 Transient Ischemic Attack (TIA) Transient ischemic attack is a temporary stroke-like event that lasts for only a short time and is caused by temporarily blocked blood vessels to part of the brain. For the purposes of HCHS/SOL classification of TIA, a TIA includes one or more episodes of [acute] focal neurologic deficit, lasting more than 30 seconds, with complete resolution of focal neurologic deficit within 24 hours. It must have no clinically relevant lesion on brain imaging (or brain imaging not done). In order for an event to be classified as TIA it must NOT have any of the following features: clonic jerking, conjugate eye deviation, prolonged focal seizure with spread, scintillating scotoma, headache with nausea and vomiting, or immediately-preceding head trauma.

A clinically relevant lesion on brain imaging includes finding judged to be consistent with signs and symptoms regardless of timing of brain imaging (i.e., less or greater than 24 hours), regardless of stroke

type (i.e., with or without blood), and regardless of imaging technique (i.e., cranial computed tomography [CT scan] or cranial magnetic resonance imaging [MRI]).

4.3.2 Stroke A stroke is defined as loss of muscle function, vision, sensation or speech resulting from brain cell damage caused by an insufficient supply of blood to part of the brain. Synonyms for stroke include apoplexy, cerebrovascular accident, or cerebral vascular accident. For the purpose of HCHS/SOL classification of stroke, a stroke involves rapid onset of neurologic deficit, headache, or meningismus and neurologic deficits not secondary to brain trauma (closed head injury), tumor, infection (e.g., encephalitis or meningitis), or other non-vascular cause. Clinical evidence or suspicion of embolic stroke secondary to SBE is counted as stroke. Classification of stroke requires either clinically relevant lesion on brain imaging*, or duration of symptoms greater than 24 hours, or death within 24 hours of symptoms.

4.4 Definitions of stroke subtypes

4.4.1 Subarachnoid hemorrhage (SAH) A subarachnoid hemorrhage has a clinical presentation of sudden onset of headache, meningismus, loss of consciousness, or coma. Focal neurologic deficit may also be present. Classification of SAH also requires consistent imaging findings with blood mainly in the subarachnoid space (basal cistern or convexity) or isolated intraventricular hemorrhage, or cerebral fluid bloody or xanthochromic on direct non-traumatic examination. Surgical or autopsy evidence of subarachnoid hemorrhage is sufficient for a classification of SAH.

4.4.2. Intraparenchymal hemorrhage (IPH) A intraparenchymal hemorrhage has a clinical presentation of focal neurologic deficit, coma as a possible accompanying condition. Classification of IPH requires consistent imaging findings with mainly intraparenchymal, dense hemorrhage, or if there is no imaging available, cerebral spinal fluid bloody or xanthochromic on direct non-traumatic examination or surgical or autopsy evidence of intraparenchymal hemorrhage is sufficient for a classification of IPH.

4.4.3. Other hemorrhage (OH) If there is insufficient data to classify subarachnoid or intraparenchymal hemorrhage, and imaging shows blood in the parenchyma, subarachnoid space, or both then a classification of other hemorrhage is made. Cerebrospinal fluid bloody or xanthochromic on direct non-traumatic examination, or surgical or autopsy evidence of blood in parenchyma, subarachnoid space, or both in the absence of a classification of SAH or IPH is sufficient for a classification of other hemorrhage.

4.4.4. Brain infarction (ischemic stroke) (INF) If a case does not meet criteria for SAH, IPH, or OH and there is a clinical presentation of focal neurologic deficit (with or without coma present) and there is consistent imaging findings without clinically relevant lesion or with clinically relevant mainly non-hemorrhagic lesion or hemorrhagic lesion indicating a hemorrhagic infarction a classification of INF can be made. Surgical or autopsy evidence of brain infarction is also sufficient for a classification of INF.

4.4.5. Other stroke types (OS) If a case does not meet criteria for SAH, IPH, OH, or INF and there is a clinical presentation of focal neurologic deficit it may be classified as OS. Examples include venous thrombosis with bleed and arterial dissection.

4.4.6. Unknown stroke type If a case does not meet criteria for SAH, IPH, OH, INF, or OS and there is a clinical presentation of focal neurologic deficit it may be classified as UNK. Examples includes evidence of symptoms but no work-up was done. A classification of NO stroke is made if the case meets none of the criteria above.

4.5 Ischemic Stroke Classification

Ischemic Stroke Classification (modified TOAST criteria utilizing findings of Neuroimaging, especially MRI and MRA) Hospitalizations classified by HCHS/SOL reviewers as ischemic strokes (INF) are further classified by reviewers using the following sub-classification criteria.

4.5.1. Cardioembolic stroke Symptoms consistent with brain infarction (see general definition of brain infarction); and acute or subacute appearing, cortical or cerebellar infarcts and brain stem or subcortical hemispheric infarcts greater than 1.5 cm in diameter on neuroimaging (CT or MRI), either in one or multiple vascular territories; and evidence suggestive for high-risk or medium-risk cardiac source for emboli (clinical and laboratory evidence); and no supportive evidence by vascular imaging for a stenosis of greater than 50% of the appropriate extracranial artery proximal to the infarct (diagnosed either by ultrasound studies, CTA, MRA, or conventional DSA); and no laboratory abnormality suggestive for non-cardiac, non-lacunar, or non-atherothromboembolic cause (for example nonatherosclerotic vasculopathies, hypercoagulable states, or hematologic disorders).

4.5.2 Extracranial large-artery atherosclerosis Symptoms consistent with brain infarction (see general definition of brain infarction); and acute or subacute appearing, cortical or cerebellar infarcts and brain stem or subcortical hemispheric infarcts greater than 1.5 cm in diameter on neuroimaging (CT or MRI); and supportive evidence by vascular imaging of a stenosis of greater than 50% of the appropriate extracranial artery proximal to the infarct (diagnosed either by ultrasound studies, CTA, MRA, or conventional DSA); and no evidence suggestive for high-risk or medium-risk cardiac source for emboli (clinical and laboratory evidence); and no laboratory abnormality suggestive for non-cardiac, non-lacunar, or non-atherothromboembolic cause (for example nonatherosclerotic vasculopathies, hypercoagulable states, or hematologic disorders).

4.5.3. Intracranial large-artery atherosclerosis Symptoms consistent with brain infarction (see general definition of brain infarction); and acute or subacute appearing, cortical or cerebellar infarcts and brain stem or subcortical hemispheric infarcts greater than 1.5 cm in diameter on neuroimaging (CT or MRI); and supportive evidence by vascular imaging of a stenosis of greater than 50% of the appropriate intracranial artery proximal to the infarct (diagnosed either by ultrasound studies, CTA, MRA, or conventional DSA); and no evidence suggestive for high-risk or medium-risk cardiac source for emboli (clinical and laboratory evidence); and no laboratory abnormality suggestive for non-cardiac, non-lacunar, or non-atherothromboembolic cause (for example nonatherosclerotic vasculopathies, hypercoagulable states, or hematologic disorders).

4.5.4. Lacunar Symptoms consistent with brain infarction (see general definition of brain infarction); and symptoms consistent with a lacunar stroke syndrome (i.e. pure motor, pure sensory, mixed sensorymotor syndrome without cortical signs, ataxic hemiparesis, dysarthria-clumsy hand syndrome); acute or subacute appearing infarct <1.5 cm in diameter in a subcortical structure or brainstem in the territory of a small penetrating artery on neuroimaging (CT or MRI): Infarct should be smaller than 1.5 cm (or described as ‘small’ or ‘lacunar’). Typical subcortical locations include corona radiata, internal capsule, external capsule, basal ganglia, thalamus. Lesions may also be located in the pons or medulla. Reviewers may score this response if ANY one of multiple new lacunar infarcts is appropriate for symptoms. Reviewers may also be scored if there is a clinically appropriate lesion of undetermined age. Do not score if the patient is described as having ‘Binswanger’s disease (encephalopathy)’ unless there is a discrete new lesion appropriate for symptoms; and no evidence suggestive for high-risk or medium-risk cardiac source for emboli (clinical and laboratory evidence); and no evidence for intra- or extracranial large artery

stenosis in the artery proximal to the infarct (by CTA, MRA, or conventional angiography); and no laboratory abnormality suggestive for non-cardiac, non-lacunar, or non-atherothromboembolic cause (for example nonatherosclerotic vasculopathies, hypercoagulable states, or hematologic disorders).

4.5.5. Other determined etiology Symptoms consistent with brain infarction (see general definition of brain infarction); and acute or subacute appearing infarct of any size in diameter either in cortical or subcortical structure or brainstem on neuroimaging (CT or MRI); and no evidence suggestive for high-risk or medium-risk cardiac source for emboli (clinical and laboratory evidence); and no evidence for intra- or extracranial large artery stenosis in the artery proximal to the infarct (by CTA, MRA, or conventional angiography); and laboratory abnormality suggestive for non-cardiac, non-lacunar, or non-atherothromboembolic cause (for example nonatherosclerotic vasculopathies, hypercoagulable states, or hematologic disorders).

4.5.6. Undetermined etiology, complete Symptoms consistent with brain infarction (see general definition of brain infarction) ; and acute or subacute appearing, cortical or cerebellar infarcts and brain stem or subcortical hemispheric infarcts greater than 1.5 cm in diameter on neuroimaging (CT or MRI); and no supportive evidence by vascular imaging of a stenosis of greater than 50% of the appropriate intra- or extracranial artery proximal to the infarct (diagnosed either by ultrasound studies, CTA, MRA, or conventional DSA); and no evidence suggestive for high-risk or medium-risk cardiac source for emboli (clinical and laboratory evidence); and no laboratory abnormality suggestive for non-cardiac, non-lacunar, or non-atherothromboembolic cause (for nonatherosclerotic vasculopathies, hypercoagulable states, or hematologic disorders).

4.5.7. Undetermined etiology, incomplete evaluation Symptoms consistent with brain infarction (see general definition of brain infarction); and incomplete cardiac evaluation (EKG or transthoracic echocardiography not performed); or incomplete evaluation of extra- or intracranial arteries (carotid ultrasound, CTA or MRA not performed).

4.5.8. Multiple possible etiologies Symptoms consistent with brain infarction (see general definition of brain infarction); and a combination of at least 2 or more of the following:

- Cortical or cerebellar lesions and brain stem or subcortical hemispheric infarcts greater than 1.5 cm in diameter on neuroimaging (CT or MRI)
- Cortical or cerebellar lesions and brain stem or subcortical hemispheric infarcts less or equal than 1.5 cm in diameter on neuroimaging (CT or MRI)
- supportive evidence by vascular imaging of a stenosis of greater than 50% of the appropriate extracranial artery proximal to the infarct (diagnosed either by ultrasound studies, CTA, MRA, or conventional DSA).
- supportive evidence by vascular imaging of a stenosis of greater than 50% of the appropriate intracranial artery proximal to the infarct (diagnosed either by ultrasound studies, CTA, MRA, or conventional DSA).
- evidence suggestive for high-risk or medium-risk cardiac source for emboli (clinical and laboratory evidence)

- no laboratory abnormality suggestive for non-cardiac, non-lacunar, or non-atherothromboembolic cause (for example nonatherosclerotic vasculopathies, hypercoagulable states, or hematologic disorders).

4.6 Ischemic Stroke Classification Use of Probable and Possible Brain Infarct Subtype Classification (modified TOAST criteria)

A "possible" diagnosis is made when the clinical findings and neuroimaging data suggest a specific subtype but other studies are not done. A "probable" diagnosis is made when the clinical findings and diagnostic work-up are complete in order to allow classification to one of the major brain infarct subtypes. The diagnostic work-up must include clinical syndrome description, neuroimaging, intra- and extracranial vascular imaging, EKG, at least transthoracic echocardiography, and basic laboratory studies.

5.0 Endpoint Surveillance for Exacerbations due to Chronic Obstructive Pulmonary Disease or Asthma

5.1 Introduction

All cases of exacerbation due to COPD or asthma resulting in either a hospitalization or emergency room visit among HCHS/SOL participants will be identified through the annual follow up call. All eligible events will be investigated and processed through the HCHS/SOL Event Classification Committee. Self-reported COPD and asthma exacerbations resulting in outpatient diagnosis and treatment without hospitalization or emergency department visits will also be identified through the annual follow up call. However, the outpatient records will not be obtained for verification.

COPD and asthma are the most common chronic lung diseases in adults. Patients with asthma have intermittent airway obstruction while those with COPD have irreversible airway obstruction (90% due to smoking). The distinction between asthma and COPD in adults has become blurred during the past five years, as the term COPD has been advertised to primary care physicians and the general public. Industry-sponsored COPD guidelines cause many adults with asthma (even never-smokers) to be falsely labeled as COPD. Asthma can begin at any age (even after age 75). Only half of asthma in adults is associated with or triggered by allergies (extrinsic asthma). There is some overlap between COPD and asthma.

By definition, GOLD criteria classify a non-smoking asthmatic that develops irreversible changes as COPD, although this type of case would be clarified clinically as COPD secondary to asthma. About half of adults with asthma are smokers. Both asthma and COPD are under-diagnosed and over-diagnosed, so self-reports and physician diagnoses are unreliable. Misclassification rates are high because objective test results are often not obtained or are incorrectly interpreted. In particular, pulmonary function tests are not generally obtained at the time of a respiratory exacerbation. About half of those with asthma or COPD in general population samples have not been diagnosed. On the other hand, some patients who have been prescribed inhalers for asthma or COPD have neither asthma nor COPD.

5.2 Event Identification

Hospitalization and emergency room events to be investigated for exacerbation due to COPD should have one of the following ICD-9 codes: 490, 491 (chronic obstructive bronchitis), 492 (emphysema), 494 (bronchiectasis), 496 (chronic airway obstruction not otherwise classified), 415, and 416.9. The codes for asthma start with 493. ICD Code 518 (Respiratory Failure, secondary to COPD) will also be investigated.

Hypersensitivity pneumonitis (495) will not be considered as COPD or asthma.


Data from hospitalizations and emergency room visits with the above discharge codes will be abstracted using the HCHS/SOL pulmonary event abstraction form (PUL). Requests for medical records are sent to the hospital and specified components of the medical record for that visit are requested. The record is blinded at the field center site and then copied and sent to the coordinating center. For emergency room visits, this information will include discharge diagnoses, physician, triage, and nurses notes, chest X-ray reports, blood count, temperature, oxygen saturation by pulse oximetry, spirometry or peak flow, arterial blood gases, and treatments such as bronchodilators, oxygen, and corticosteroids. For hospitalizations, the same test results will be required, and in addition, discharge medications and pulmonary function results, if performed.

Exacerbations of both asthma and COPD will be diagnosed based on chart review including the materials described above. In a patient presenting to the emergency department (ED) with an acute episode of shortness of breath, it is often difficult to distinguish between pneumonia, an asthma exacerbation, a COPD exacerbation, a pulmonary embolism, and an exacerbation of congestive heart failure (CHF). A bacterial or viral pneumonia can trigger an asthma or COPD exacerbation, but the usual cause is an upper respiratory viral infection (Fabbri L, et al, Similarities and discrepancies between exacerbations of asthma and COPD. *Thorax* 1998; 53:803). The criteria used below were created by consensus by the HCHS/SOL pulmonary endpoints committee.

5.3 Diagnostic criteria for pulmonary events

Initially reviewers will state the level of evidence for whether the participant has a *history of chronic lower respiratory disease* (CLRD) as definite, probable, probably not, definitely not, and unknown. CLRD includes COPD, emphysema, chronic bronchitis, and asthma. Our first question is broad, in case details regarding history of respiratory disease are limited. Criteria for definite and probable CLRD are shown in the figure below. Criteria for a categorization of “definite” for each type of CLRD are specified below in the figure.

Figure 2.



The peak expiratory flow (PEF) and pulmonary function tests (PFTs) referred to in the figure are those in the medical record and not from the HCHS/SOL study visit. PFTs from the study visit will not be used to classify events. *See PFT criteria page for definitions of obstruction.

Definite criteria for the diagnosis of COPD require obstruction on PFTs or a low peak expiratory flow, both defined in the PFTs section. Definite emphysema must be confirmed on chest CT. Definite chronic bronchitis is only a clinical definition: cough for at least 3 months of the year for at least two years in a row. Note that you can have definite emphysema without PFTs, but this would not be definite COPD (need to confirm obstruction). The same is true for chronic bronchitis (ie, no PFTs does not confirm COPD)

Definite asthma is defined by low peak flow with improvement with bronchodilators in a patient less than 45 years of age. Definite criteria for diagnosis of asthma also includes the following: positive methacholine challenge test, and reversibility on PFTs with no other obstructive lung diseases (eg CF, bronchiectasis, upper airway obstruction). See below for definition of reversibility in PFTs. In hospital and ER medical records are most likely to have PEF measures and not PFT measures.

‘PROBABLE’ CLRD should be selected if there is a physician diagnosis of COPD, emphysema, asthma, or chronic bronchitis OR chronic use of medications used to manage CLRD including inhaled corticosteroids (not intra-nasal), inhaled long acting beta agonists (eg, salmeterol [Serevent], formoterol [Foradil]), inhaled long acting anticholinergic (ie Spiriva or tiotropium), or methylxanthines (eg, theophylline).

If the answer is definite or probable history of CLRD, we ask about the evidence for history of each specific type of CLRD --COPD, emphysema, chronic bronchitis, and asthma with the same categories as in the first question. These questions go beyond ‘history of’, include ‘new onset’.

COPD. Answer DEFINITE if there is definite or probable history of CLRD and if there is evidence of obstruction from PFTs, per HCHS/SOL criteria. IF PFTs are not available and there is a history reported of COPD then answer PROBABLE. IF there is conflicting information as to whether the patient has COPD or not then answer unclassifiable. If there are normal PFTs then answer DEFINITELY NOT.

Emphysema. Answer DEFINITE if there is evidence of emphysema from a CT scan. Answer PROBABLE if there is an MD diagnosis of emphysema, but there is not imaging to support the diagnosis or if only the CXR is suggestive of emphysema. Answer DEFINITELY NOT, if there is a normal CHEST CT in which the lung parenchyma is reported as normal. IF there is conflicting information as to whether the patient has emphysema or not then answer unclassifiable.

Chronic Bronchitis. Definite chronic bronchitis is only a clinical definition. Answer DEFINITE if cough for at least 3 months of the year for at least two years in a row is reported. Answer PROBABLE, if there is a physician diagnosis of chronic bronchitis. Answer UNCLASSIFIABLE if it is not stated whether the patient has had chronic cough or not.

Asthma. Answer DEFINITE, if patient with a history of asthma has a low peak flow with improvement with bronchodilators AND the patient is less than 45 years of age. Definite criteria for diagnosis of asthma also includes the following: positive methacholine challenge test, and reversibility on PFTs with no other obstructive lung diseases (eg CF, bronchiectasis, upper airway obstruction). In medical records, we are most likely to have PEF measures. Answer PROBABLE, if patient has a history of asthma AND has a history of the following: age of onset of asthma under the age of 40, positive allergy testing, wheezing on exam that responds to bronchodilators.


If there is some other lung disease present, then specify the type.

Next reviewers are asked about an exacerbation of underlying chronic lower respiratory disease (CLRD). Answer PROBABLE, if the patient has new onset or worsening of respiratory symptoms which must include the following symptoms: dyspnea, cough, change in sputum color, change in sputum volume, or wheeze (either reported or on exam). If the primary reason the patient is in the ER or hospital is for one of these symptoms then assume symptoms are worse from baseline. Answer 3a.

Answer DEFINITE, if the patient has criteria for a probable CLRD exacerbation AND a low peak flow or low FEV1/FVC, AND absence of alternative diagnosis likely to cause CLRD symptoms and/or low peak flow or low FEV1/FVC. Answer DEFINITELY NOT, if the patient does not have symptoms of CLRD exacerbation and/or has a clear alternative cause of symptoms. Answer PROBABLY NOT, if there is not strong enough evidence to answer DEFINITELY NOT, but you feel it is not likely that the patient has an exacerbation of CLRD. Answer UNCLASSIFIABLE if there is inadequate information to determine if there is an exacerbation of CLRD. Skip to question 4.

Figure 3.

HCHS/SOL criteria for Exacerbation of CLRD



If definite or probable exacerbation, then specify which type of lung disease is likely the cause: COPD predominant, asthma predominant, either asthma or COPD or unclassifiable. Answer 'asthma predominant' if it is clear from the medical record that asthma is the cause of the exacerbation. Answer 'COPD predominant' if it is clear from the medical record that COPD is the cause of the exacerbation. Answer 'either asthma or COPD' if the physicians are not sure if the patient has asthma or COPD and/or if there is conflicting evidence as to which diagnosis that is the cause of the exacerbation. Answer UNCLASSIFIABLE if there is not adequate information to determine whether COPD or asthma was the cause of the exacerbation.

Pneumonia causes respiratory symptoms that are similar to a COPD exacerbation, it is somewhat controversial as to whether person with COPD and pneumonia also has a COPD exacerbation, therefore

reviewers need to state whether the patient had pneumonia, defined as a new infiltrate on Chest Xray or on chest CT scan. The same 5 categories from definite to unknown are available as options.

Interpretation of PFTS and Peak expiratory flow from medical records

In order to classify pulmonary events, one must have criteria for interpreting pulmonary function tests and peak expiratory flow values that are found in the medical record. Although as mentioned, it is mostly PEF values that will be found in a medical record for a potential CLRD exacerbation. The criteria described and summarized below.

Interpretation for Pulmonary Function Tests and Peak Expiratory Flow (PEF) from Medical Records

- “Low PEF”
 - PEF < 70% predicted*
- Low PEF “w/ improvement”
 - Increase in PEF of $\geq 30\%$ or, in absence of repeat PEF, clear improvement in clinical status
- “Low FEV₁/FVC ratio”
 - Pre-bronchodilator FEV₁/FVC < LLN[†]
- “COPD”
 - Post-bronchodilator FEV₁/FVC ratio < 0.70 or < LLN[†] AND FEV₁% predicted < 80%[†]
 - In absence of post-bronchodilator measures, FEV₁/FVC ratio < 0.70 or < LLN[†] AND FEV₁% predicted < 65%[†]

“Low PEF” is defined as PEF < 70% predicted. PEF predicted values will be based upon HCHS/SOL predicted values (currently pending) using age at time of clinical PEF measurement, gender, and measured height at HCHS/SOL exam. These 3 variables are included in the event summary form available with all medical records sent to reviewers.

Low PEF “w/ improvement” is defined as an increase in PEF of $\geq 30\%$ or, in absence of repeat PEF, clear improvement in clinical status.

“Low FEV₁/FVC ratio” is defined as pre-bronchodilator FEV₁/FVC < LLN (lower limit of normal). The LLN for FEV₁/FVC is based upon HCHS/SOL reference equations, determined on each individual patient, (currently pending) using age at time of clinical PFT measurement, gender, and measured height at HCHS/SOL exam.

“COPD” from PFT results are based ideally on post-bronchodilator PFTS; however, if not available then pre-bronchodilator PFTs are used.

The definition for COPD from post-bronchodilator PFTs is: 1) FEV₁/FVC ratio < 0.70 or < LLN.

In absence of post-bronchodilator measures, the definition of COPD from pre-bronchodilator PFTS is: 1) FEV1/FVC ratio <0.70 or <LLN AND 2) FEV1 percent predicted <65%.

6.0 Endpoint Surveillance for Pregnancy Related Complications

6.1 Introduction

The numerous physiological changes that occur during pregnancy stress the metabolic system and thus may reveal existing subclinical disease states as well as trigger the onset of new chronic conditions. Several studies have demonstrated that gestational diabetes (GDM) is associated with a four-fold higher risk of subsequent type 2 diabetes (T2DM). Hypertensive disorders during pregnancy are associated with subsequent high CVD risk factor status and elevated CVD risk. Offspring born to women with GDM are more likely to be overweight or develop metabolic syndrome or T2DM later in life. However, few such studies have included H/L women, and few studies have pre-pregnancy metabolic measures from a population-based sample. HCHS/SOL allows timely study of this issue, given the high prevalence of obesity (a RF for pregnancy complications) among H/L women, and the proposed collection of extensive longitudinal data on CVD RFs.

The specific endpoints of interest for pregnancy related complications (PRC) are pre-eclampsia, eclampsia, and gestational diabetes. All potential cases of one of these pregnancy related complications resulting in either a hospitalization or emergency room visit among HCHS/SOL participants will be identified through the annual follow up call, at the time of the second HCHS/SOL study visit, or through existing hospital medical records available at the CSCC. All eligible events will be investigated and processed through the HCHS/SOL Event Classification Committee. In addition, if possible, birth certificates will be requested to provide information on the child, and assure other events were not missed. Note that these three conditions will be classified by reviewing all hospitalizations or ER visit, and the birth certificate (if available), for the entire pregnancy episode, rather than by each hospitalization or ER visit.

6.2 Event Identification

Hospitalization and emergency room events to be investigated for pregnancy related complications (PRC) of interest should have one of the relevant keywords (see table) in the medical record for a pregnancy or one of the following ICD-9 or ICD-10 codes shown in the table. Since baseline, the HCHS/SOL study has collected discharge summaries and face sheets for all hospitalizations. Those hospital medical records on file at the CSCC will be used to classify PRC using an abstraction form, and a physician review panel with adjudication.

Medical records collected for all hospitalizations since baseline will be used to classify pregnancy related complications events using those existing records. In addition, we will request birth certificates for all births since the baseline visit to assess the presence of maternal pregnancy-related diagnoses of interest for each pregnancy. In addition, the event eligibility form has been updated to include the relevant keywords and ICD codes for PRC, such that current hospitalizations and ER visits that are being investigated will request additional medical records of relevance to classify the event.

Data from hospitalizations and emergency room visits with any of the below discharge codes or keywords will be abstracted using the HCHS/SOL pregnancy complications abstraction form (PCA). Requests for medical records are sent to the hospital and specified components of the medical record for that visit are requested. The record is blinded at the field center site and then copied and sent to the coordinating center.

Pregnancy related complications will be diagnosed per pregnancy rather than per admission or ER visit. All admissions and emergency room visits for that pregnancy will be reviewed as a single event by physicians on the pregnancy related complications review panel. Events for pregnancy related complications will be diagnosed based on chart review including the materials described above.

Table 6. Keywords and ICD codes used to identify events for abstraction as and review as a possible pregnancy related complication			
Diagnosis	ICD9	ICD10	Keywords
Mild or unspecified preeclampsia	642.4x	O14.0x, O14.9x	Preeclampsia, toxemia, eclampsia, PreX, PEC, PET,
Severe preeclampsia	642.5x	O14.1x O14.2x	Preeclampsia, toxemia, eclampsia, PreX, PEC, PET,
Preeclampsia superimposed on chronic HTN	642.7x	O11.x	Preeclampsia, toxemia, eclampsia, PreX, PEC, PET,
Gestational hypertension	642.3x	O13.x	PIH, pregnancy-induced hypertension, gHTN, Gestational hypertension
Eclampsia	642.6x	O15.x	Seizure, eclampsia,
Pregestational hypertension	642.0x 642.1x 642.2x	O10.1x O10.2x O10.3x O16.x	Chronic hypertension, pregestational hypertension, htn
Gestational Diabetes	648.8x	O24.4 O99.81	Gestational diabetes, diabetes, hyperglycemia, insulin
Pregestational Diabetes	648.0x	O24.0x O24.1x O24.3x O24.8x O24.9	Gestational diabetes, diabetes, hyperglycemia, insulin

6.3 Physician review of events

There will be a panel of physician reviewers for pregnancy related outcomes. This panel will be trained and certified to classify PRC. There will be one type of reviewer form for all PRC of interest including pre-eclampsia, eclampsia and gestational diabetes. For each potential pregnancy event, two reviewers will be assigned to review the event independently, and if needed, an adjudicator will resolve differences between reviewers to provide a final classification for the event.

6.4 Diagnostic criteria for pregnancy related complications

There are two main outcomes of interest for pregnancy related complications, which are: 1) Pre-eclampsia or eclampsia and 2) Gestational Diabetes. The diagnostic criteria that will be used to classify these events by the physician reviewer panel are discussed below. Criteria for each of these pregnancy complications

will be defined based on existing diagnostic criteria, and experience from research studies. Physician reviewers are asked to use their judgment in applying these criteria after review of the available medical records.

6.4.1 Pre-eclampsia and Eclampsia

Table 7. Diagnostic criteria for definite, probable, unlikely and unclassifiable pre-eclampsia and eclampsia		
	Pre-Eclampsia	Eclampsia
Definite	1) Pregnancy-related hypertension (SBP ≥ 140 and DBP ≥ 90 on two occasions at least 6 hours apart) AND 2) Proteinuria by one of the following criteria one of the following: a) $\geq 0.3g / 24h$ urine , or b) $\geq 1+$ urine dipstick, or c) urine protein:creatinine ratio documented as “positive” by local lab criteria	1) Pregnancy-related hypertension (SBP ≥ 140 and DBP ≥ 90 on two occasions at least 6 hours apart) AND 2) a grand mal seizure during pregnancy in the absence of other known causes
Probable	1) a) Mention of pre-eclampsia, or other keywords for pre-eclampsia in the medical record AND b) Mention of protein or albumin in the urine in the medical record OR 2) Documentation of intrapartum magnesium for seizure prophylaxis	A grand mal seizure during pregnancy in the absence of other known causes for seizure
Unlikely	Normal documented blood pressure, AND urine protein not documented or $<1+$, AND no mention of preeclampsia for this pregnancy in obstetric records in a record with reasonable documentation	No mention of seizure or eclampsia during this pregnancy, in a medical record with reasonable documentation
Unclassifiable	Contradictory information or inadequate medical record such that unable to classify as pre-eclampsia or not	Contradictory information or inadequate medical record such that unable to classify as eclampsia or not

6.4.2 Gestational Diabetes Mellitus

Definite	<p>1) Glucose tolerance testing results not available AND documented treatment with insulin or oral hypoglycemic (ie glyburide) OR</p> <p>2) Laboratory evidence of diagnostic glucose intolerance (one of a-d):</p> <p style="padding-left: 40px;">a) 3 hour Oral glucose tolerance test (OGTT) with ≥ 2 abnormal values: fasting glucose ≥ 95mg/dL; 1 hour ≥ 180mg/dL; 2 hour ≥ 155mg/dL; 3 hour ≥ 140mg /dL (Carpenter-Coustan Criteria), OR</p> <p style="padding-left: 40px;">b) 2 hour 75g oral glucose tolerance test exceeding any one of the following thresholds: fasting glucose ≥ 92 mg/dl, or 1 hour ≥ 180 mg/dl, or 2 hour ≥ 153 mg/dl (IADPSG criteria), OR</p> <p style="padding-left: 40px;">c) 50g GLT exceeding 200 mg/dl, OR</p> <p style="padding-left: 40px;">d) Two or more documented fasting glucoses > 125 mg/dL</p>
Probable	1) Physician or advanced practice provider mention of current gestational diabetes in medical record
Possible	<p>1) Laboratory evidence of abnormal but non-diagnostic glucose intolerance:</p> <p style="padding-left: 40px;">a) Elevated 1-hour Glucose load testing ≥ 135 mg/dl, OR</p> <p style="padding-left: 40px;">b) Other evidence of abnormal but non-diagnostic glucose tolerance: fasting glucose ≥ 105 mg/dl, or 1 hour post prandial ≥ 140 mg/dl, or 2 hour post prandial ≥ 120 mg/dl)</p>
Unlikely	No evidence in medical record to suggest deviation from normal glucose tolerance
Unclassifiable	Contradictory information or inadequate medical record such that unable to classify as GDM or not

7.0 Endpoint Surveillance of Fatal Events

7.1 Event Identification

Deaths will be identified using several methods including questions asked during the annual follow up contact, review of vital statistics lists and obituaries from each state, or matching with the National Death Index (NDI). Once a death is identified, field center staff will obtain a death certificate and send a copy to the coordinating center for processing and abstraction. Deaths requiring investigation will be those with the following codes:

ICD-10 codes (Underlying cause of death)	Description
I00-I52, I70-I99	Cardiovascular Disease
I60-I69	Cerebrovascular Disease
G45-G46	Transient cerebral ischemic attack
E10-E14	Diabetes
J81	Pulmonary edema
R96, R98, R99	Ill-defined
R07	Chest Pain
J40-J42	Bronchitis
J43	Emphysema
J44	Other chronic obstructive pulmonary disease
J45	Asthma
J46	Status asthmaticus (acute severe asthma)
J47	Bronchiectasis

7.2 Diagnostic Criteria

7.3 Fatal Coronary Heart Disease Events

The 2003 Scientific Statement (Luepker, 2003) recommended the following case classification of fatal CHD events for hospitalized patients:

- A. Definite fatal MI
 1. Death within 28 days of hospital admission in definite MI cases
 2. Postmortem findings consistent with MI within 28 days
- B. Probable fatal MI
 1. Death within 28 days of hospital admission in cases defined in probable MI cases
 2. Death within 6 hours of hospital admission with cardiac symptoms and/or signs. Other confirmatory data (biomarkers, ECG) are absent or not diagnostic.
- C. Possible fatal coronary event
 1. Death within 28 days of hospital admission for possible MI, unstable angina or chronic stable angina
 2. Postmortem findings show old infarct and/or $\geq 50\%$ atherosclerotic narrowing of coronary arteries.

Out-of-hospital CVD death is a major public health burden, accounting for 50-75% of all fatal CVD events in countries in which it is documented. Classifying out-of-hospital deaths will be a part of the HCHS/SOL outcomes and adjudication activities. Classification of out-of-hospital CHD death is usually deficient because of its sudden onset, lack of information from the victim, lack of witnesses, and low autopsy rates.

Information recommended by the Scientific Statement for adjudication of all deaths, including out-of-hospital deaths, will include:

1. Site of death
 - A. Before transport to medical facility (e.g., home, work site, street, nursing home)
 - B. During transport (e.g., ambulance, car)
 - C. Pronounced dead in emergency department and not admitted to hospital
2. Cause
 - A. Death certification, based on WHO methods for assigning causation, is the primary source
 1. Validated by postmortem examination, if available
 2. Enhanced by other information, if available (see 2.B and C below)
 - B. Interviews with witnesses and family members
 - C. Medical history from healthcare records and physicians
3. Timing
 - A. One hour and/or 24 hours since last seen or known to be alive
 - B. Other time categories based on research needs

HCHS/SOL staff members perform a brief interview with either next-of-kin or contacts provided by the participant for this purpose. This interview will gather information essential for adjudication. Elements will include:

1. Where did the death occur? (out of hospital, at personal residence, emergency department, etc.)
2. Was the person stable prior to death?
3. Was death sudden and unexpected? (definitely, possibly, no)
4. Was death witnessed? (If not, were they found in bed or a chair?)
5. Time from onset of symptoms (or last seen) to death, in hours (0-1, 1-6, 7-24, 24, unknown)
6. Was there a hospitalization or emergency department admission either immediately before death, or since the last follow-up contact?
7. Can informant send a death certificate including cause of death?

The 2003 Scientific Statement (Luepker, 2003) recommended a classification scheme for cause of out-of-hospital CHD death, in the following hierarchy:

- I. Definite fatal MI:
Documented definite or probable MI in the previous 28 days and no evidence of a noncoronary cause of death, or autopsy evidence of recent coronary occlusion or MI <28 days old.
- II. Definite fatal CHD:
 - (1) A history of CHD and/or documented cardiac pain within 72 hours before death and
 - (2) no evidence of a non-coronary cause of deathor
 - (3) autopsy evidence of chronic CHD, including coronary atherosclerosis and myocardial scarring.
- III. Possible fatal CHD:
An ICD code (underlying cause) for CHD death (ICD 9*: 410 to 414, 427.5, 429.2; ICD 10: I20 to I25 and I46) and no evidence of a non-coronary cause of death
- IV. Cardiac death:
When death certificates are the only source of information: ICD 9: 390 to 398, 402, 404 to 429; ICD 10: I00 to I09, I11, I13, I20 to I25, I27, I30 to I52

- V. Non-CHD death:
Evidence of a noncoronary cause of death
- VI. Unclassifiable:
Insufficient information to determine whether the death was a CHD death (at any certainty level) or a noncardiac death: ICD 9 code 799

7.4 Fatal Pulmonary Events

The hospital records will be obtained when a respiratory death (J42, J44, J45, J46) occurred during a hospitalization, or when an out-of-hospital death occurred within 30 days following a hospitalization. Ascertainment of cause-specific mortality in patients with COPD is difficult, but should be attempted by a central committee, not site investigators (McGarvey LP, et al. Ascertainment of cause-specific mortality in COPD: operations of the TORCH Clinical Endpoint Committee. *Thorax* 2007; 63:411).

Death certificates will be obtained in all cases. If a death certificate cannot be obtained, it will be stated in the documentation. If medical records are inadequate and a death certificate cannot be obtained, a cause of death will be adjudicated based on the best available evidence of record. If a probable cause cannot be adjudicated, it will be classified as “unknown”.

The primary cause of death should be attributed to the disorder that causes the patient to present for medical treatment and should be distinguished from terminal events that are the immediate cause of death.

For example, if a patient is admitted to the hospital with a COPD exacerbation and during the exacerbation subsequently develops complications such as pneumonia, respiratory failure, renal failure, sepsis or myocardial infarction, the primary cause of death should be attributed to COPD.

8.0 Event Classification Committee

8.1 Introduction

The Event Classification Committee (ECC) is organized into three working groups consisting of a cardiac group, a stroke group, and a pulmonary group. See Figure 1. The cardiac group is responsible for reviewing hospitalized myocardial infarction, heart failure, hospitalized cardiac death, and out of hospital cardiac death. The stroke group is responsible for hospitalized stroke and TIA reviews. The pulmonary group is responsible for reviewing hospitalized COPD and asthma as well as emergency department only events for exacerbations of COPD and asthma. A chair or lead reviewer is designated for each group.


8.2 Review Process

The Coordinating Center assembles and disseminates all review packets to the appropriate Physician Reviewers through an online system accessible through the HCHS/SOL website. The reviewers have 2 weeks, from when notified that a set of cases is ready for review, to complete a set of reviews. A reviewer’s classification of an event applies only to the specific hospitalization or emergency department visit or outpatient event under review. The reviewer should not be concerned if there is a history of prior events or records. Each event should be judged separately. Each case is reviewed independently by 2 reviewers. Disagreements will be resolved by the chair of the respective review group.

8.3 Disagreement Resolution

Cases that produce a disagreement among the two independent reviews will be sent to the chair of the appropriate review group for resolution. The chair will have access to results of each reviewer’s decisions and will make a third and final classification for these cases.

Figure 4.



8.4 Confidentiality

Several procedures are in place to protect the security of the personal identifying information obtained from medical records and used in the event ascertainment process. Personal identifying information (name, SSN, date of birth, etc) from cohort members is used for the purpose of linkage to the National Death Index. This information is needed to determine vital status of cohort participants who are lost to follow up. All personal identifiers of cohort participants', treating physicians, hospital name and location, and other identifying information are blacked out on any paper copy of medical records sent to the coordinating center and doubled checked prior to blinding for distribution to the ECC. Any copies of these documents remaining at the coordinating center are stored in locked secure rooms. ECC members are instructed in the proper confidential destruction of any medical record information they are provided. Study personnel involved in processing the medical record information, from abstraction to handling of these data have been trained on the protection of human subjects in research.

References

Luepker RV, Apple FS, Christenson RH, Crow RS, Fortmann SP, Goff D, Goldberg RJ, Hand MM, Jaffe AS, Julian DG, Levy D, Manolio T, Mendis S, Mensah G, Pajak A, Prineas RJ, et al. Case definitions for acute coronary heart disease in epidemiology and clinical research studies. A statement from the AHA Council on Epidemiology and Prevention; AHA Statistics Committee, World Heart Federation Council on Epidemiology and Prevention; the European Society of Cardiology Working Group on Epidemiology and Prevention; Centers for Disease Control and Prevention; and the National Heart, Lung and Blood Institute. *Circulation* 2003;108:2543-2549.

APPENDIX A. Minnesota Code ECG Criteria

APPENDIX B. GLOSSARY of Key Event Data Collections Terms

APPENDIX C. Data Collection Forms required for assessing Endpoints:

HCHS/SOL Event Tracking Form (ETR) & QxQ

HCHS/SOL Informant Interview Form (IIE) & QxQ

HCHS/SOL Death Certificate Form (DTH) & QxQ

APPENDIX D. Abstraction Forms

Myocardial Infarction Abstraction Form (MIF)

Heart Failure Abstraction Form (HTF)

Pulmonary Abstraction Form (PUL)

APPENDIX E. Reviewer Forms for classification and adjudication

Myocardial Infarction Form (MIF)

Stroke and TIA Form

Heart Failure Exacerbations Form (HFT)

COPD and Asthma Form (PUL)

APPENDIX F. Event Summary Forms (ESF)

Myocardial Infarction Abstraction Form (MIF)

Heart Failure Abstraction Form (HTF)

Pulmonary Abstraction Form (PUL)

APPENDIX A. Minnesota Code ECG Criteria

Table 1. Evolving diagnostic ECG (any of the following Q1 through Q4)

Evolving Q1: No Q-code in prior study ECG or first ECG in event set of ECG(s) followed by a record with a diagnostic Q-code (Minnesota Code 1-1-1 through 1-2-5 plus 1.2-7) OR any code 1-3-X or in baseline ECG followed by a record with any code 1-1-X.

Evolving Q2: An equivocal Q-code (any 1-3 code) and no major ST-segment depression in prior study ECG or first ECG in event set of ECG(s) followed by a record with a diagnostic Q-code PLUS a major ST-segment depression (Minnesota code 4-1-X or 4-2) and 100% increase in ST depression

Evolving Q3: An equivocal Q-code (any 1-3 code) and no major ST-segment depression in prior study ECG or first ECG in event set of ECG(s) followed by a record with a diagnostic Q-code PLUS a major T-wave inversion (Minnesota Code 5-1 or 5-2) and 100% increase in T-wave inversion

Evolving Q4: An equivocal Q-code and no ST-segment elevation in prior study ECG or first ECG in event set of ECG(s) followed by a record with a diagnostic Q-code PLUS ST-segment elevation (Minnesota code 9-2) and 100% increase in STE

*Note for Table 2: A significant Q-code change requires $\geq 50\%$ increase in event Q/R ratio or ≥ 1 mm initial R-wave amplitude decrease in event ECG compared with corresponding lead(s) of baseline ECG.

Table 2. Positive ECG

(a) Evolving ST elevation alone

Evolving STE 1: No 9-2 in prior ECG or first ECG in event set of ECG(s) and 9-2 in at least 2 leads of a following event ECG with 100% increase in STE in both leads.

Evolving STE 2: 9-2 in prior ECG or first ECG in event set of ECG(s) with a 100% increase in STE in at least 2 leads.

Evolving STE 3: 9-2 and no 5-1 or 5-2 in prior ECG in first ECG in event set of ECGs and appearance of 5-1 or 5-2 with 100% increase in T-wave inversions in at least 2 leads.

Evolving STE R1: Reversal of evolving STE 1.

Evolving STE R2: Reversal of evolving STE 2.

OR

b) Evolving equivocal Q-wave plus evolving ST-T depression/inversion

Evolving Q5: No Q-code and neither 4-1-X nor 4-2 in prior study ECG or first ECG in event set of ECG(s) followed by a record with an equivocal Q-code PLUS 4-1-X or 4-2 and 100% increase in ST depression

Evolving Q6: No Q-code and neither 5-1 or 5-2 in prior study ECG or first ECG in event set of ECG(s) followed by a record with an equivocal Q-code PLUS a 5-1 or 5-2 100% increase in T-wave inversion

Evolving Q7: No Q-code and no 9-2 in prior study ECG or first ECG in event set of ECG(s) followed by a record with an equivocal Q-code PLUS a 9-2 and a 100% increase in STE.

OR

c) New left bundle branch block (code 7-1-1, with the QRS duration increasing by at least 20 ms from less than 120 ms to ≥ 120 ms).

*STE indicates ST elevation. In the case of a single ECG available only from a possible event hospital admission, a probable MI can be classified if compared with the previous study ECG, there is a new appearance of a diagnostic Q-wave (+MS 1-1-1 through 1-2-5 plus 1-2-7 or any code 1-3-X in the previous ECG and the event ECG has any code 1-1-X or presence of 9-2 in at least 2 leads).

Table 3. Nonspecific ECG: Evolution of minor ST-T depression/inversion alone, or minor Q-wave alone

(a) Evolving non-STE non-Q-wave pattern MI

Evolving ST-T1: Either 4-0 (no 4-code), 4-4, or 4-3 in previous ECG or first ECG in event set of ECG(s) followed by a record with 4-2 or 4-1-2 or 4-1-1 and 100% increase in ST segment depression

Evolving ST-T2: Either 4-2 or 4-1-2 in previous ECG or first ECG in event set of ECG(s) followed by a record with 4-1-1 and 100% increase in ST segment depression

Evolving ST-T3: Either 5-0, 5-4, or 5-3 in previous ECG or first ECG in event set of ECG(s) followed by a record with 5-2 or 5-1 and 100% increase in T-wave inversion

Evolving ST-T4: Code 5-2 in previous ECG or first ECG in event set of ECG(s) followed by a record with 5-1 and 100% increase in T-wave inversion

Evolving ST-T5: Code 4-1-1 in previous ECG or first ECG in event set of ECG(s) followed by a record with 4-1-1 and 100% increase in ST depression

Evolving ST-T6: Code 5-1 in previous ECG or first ECG in event set of ECG(s) followed by a record with 5-1 with 100% increase in T-wave inversion

Evolving ST-T7: Code 5-2 in previous ECG or first ECG in event set of ECG(s) followed by a record with 5-2 with 100% increase in T-wave inversion

Evolving ST-T R1 through ST-T R7_the reverse of ST-T1 to ST-T7, respectively

OR

(b) Evolving minor Q wave alone

No Q code in previous study ECG or event ECG, followed by an event ECG with an equivocal Q-code (any 1-3 code)

The ECG series is assigned the highest category for which criteria are met, i.e., evolving diagnostic > diagnostic > evolving ST-T patterns > equivocal > other.

APPENDIX B. GLOSSARY of Key Event Data Collections Terms

This section has intentionally been left blank.

APPENDIX C. Data Collection Forms required for assessing Endpoints:

- HCHS/SOL Event Tracking Form (ETR)
- HCHS/SOL Informant Interview Form (IIE)
- HCHS/SOL Physician Interview Form (PQE)
- HCHS/SOL Death Certificate Form (DTH)

Note that current copies of all study data collection forms and QxQs can also be found on the study web site page:

http://www.csc.unc.edu/hchs/utilities/docfilter.php?study=hchs&filter_type=forms



HCHS / SOL Event Tracking form

PARTICIPANT ID NUMBER:											
------------------------	--	--	--	--	--	--	--	--	--	--	--

FORM CODE:
ETR
VERSION: A
07/01/09

Contact Occasion			SEQ #		
------------------	--	--	-------	--	--

EVENT ID NUMBER:

EVENT Date: / /

Instructions: This form is completed by the Outcomes Coordinators to document the stages in the hospital or Emergency Department records acquisition process. The EVENT ID and DATE of EVENT fields above are pre-filled by the Data Management System for HCHS. The form is entered into the DMS as a multi-line form with the last status being the one of record for a given event. Use as many paper forms as needed to track the progress of medical records acquisition.

Event Records Tracking Results

1. Date (MM/DD/YYYY)	2. Notes	3. Result Code*	4. Staff Code
/ /			
/ /			
/ /			
/ /			
/ /			
/ /			
/ /			

*RESULT CODES for Records Processing

- 0 Pending records request
- 1 Release of Information requested from Participant
- 2 Release of Information obtained from Participant
- 3 Event Record requested
- 4 Confirmed, No event to investigate
- 5 Confirmed, Records Not Available
- 6 Medical records received for event
- 7 Supplemental records requested
- 8 Verification of medical records to be sent [note: ICD-9 codes needed at this step]
- 9 Shipping medical records to Coordinating Center [cover sheet for shipping produced]



HCHS/SOL Informant Interview

ID NUMBER:	<input type="text"/>								
-------------------	----------------------	----------------------	----------------------	----------------------	----------------------	----------------------	----------------------	----------------------	----------------------

FORM CODE: IIE
VERSION: A
12/02/2008

Contact Occasion	0	1	SEQ #	<input type="text"/>	<input type="text"/>
------------------	---	---	-------	----------------------	----------------------

Administrative Information

0a. Completion Date: / /
Month Day Year

0b. Staff ID:

Public reporting burden for this collection of information is estimated to average 30 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to, a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to: NIH, Project Clearance Branch, 6705 Rockledge Drive, MSC 7974, Bethesda, MD 20892-7974, ATTN: PRA (0925-0584). Do not return the completed form to this address.

Instructions: The informant interview form is completed for each informant for an eligible death as determined by the HCHS/SOL event investigation worksheet.

<p>Decedent's name: _____ _____</p> <p>Date of death: <input type="text"/><input type="text"/>/ <input type="text"/><input type="text"/>/ <input type="text"/><input type="text"/><input type="text"/><input type="text"/></p> <p>Age at death: <input type="text"/><input type="text"/></p> <p>Place of death: _____</p>	<p style="text-align: center;">Informant name:</p>
---	--

"Hello, my name is (interviewer's name) with the HCHS/SOL study. I'm calling regarding (name of decedent) involvement with the HCHS/SOL study, a medical study in which (name of decedent) was enrolled. I want to express our condolences for your loss. We understand that you have been identified as someone who can help us close out (decedent name)'s file. I need to ask you a few questions about the circumstances surrounding (name)'s death. Would now be a good time to talk?"

No —————> When would be convenient to call back?

Yes —————> Thank you. If you have any questions, please ask me.

1. "Before we get started could you please tell me what was your relationship with the deceased?" (Respondent was deceased's...)

- Spouse 1
- Daughter/Son 2
- Parent 3
- Friend 4
- Workmate 5
- Other relative 6 Specify: _____
- Other 7 Specify: _____

“Now, I would like to ask you about the circumstances surrounding (insert decedent’s name) medical history.”

2. *“Please tell me about his/her general health, health on the day s/he died, and about the death itself.”*

Record a brief synopsis of the events surrounding the death as related by the informant:

“Some of the remaining questions may repeat information already provided, but it helps us to ask these items specifically.”

3a. Was anyone present when s/he died?

- No 0
Yes 1
Unknown 9

3b Where was (insert decedent name) when s/he died (Check one)

- Home 0 → **GO TO 3c**
Work 1
Public building 2
Bus or public transportation 3
In a car 4
Nursing home 5
In an emergency room 6
In an ambulance 7
In a hospital 8
Unknown 9
Other 10

3c. If s/he died at home, was s/he found:

- In bed 0
In a chair 1
On the floor 2
Elsewhere 3 specify: _____
Unknown 9

4. Was anyone close enough to hear (insert decedent's name) if s/he had called out?

- No 0
- Yes 1
- Unknown 9

5. How long was it between the time (insert decedent's name) was last known to be alive and the time s/he was found dead?

- Less than 5 minutes 1
- 5 minutes to 1 hour 2
- 1 to 24 hours 3
- Longer than 24 hours 4
- Unknown 9

6. Please tell me who was present. (check all that apply)

- Self 1 (skip to question 8)
- Health care person(s) 2
- Other person(s) 3

7. When was the last time you saw (insert decedent's name) prior to his/her death?

- Less than 5 minutes 1
- 5 minutes to 1 hour 2
- 1 to 24 hours 3
- Longer than 24 hours 4
- Unknown 9

HISTORY

The next few questions concern (insert decedent's name) medical history.

8. Was s/he restricted to home, able to leave home only with assistance or great effort, or was his/her activity unrestricted?

- Restricted to home 1
- Able to leave home only with assistance or great effort 2
- Unrestricted 3

9. Was s/he hospitalized within the four weeks prior to death?

- No 0 Skip to question 13
- Yes 1
- Unknown 9 Skip to question 13

10. What was the reason for the hospitalization? (select all that apply)

- Heart attack or heart disease 1
- Stroke 2
- Heart surgery 3
- Surgical procedure (other than heart) 4
- Emphysema, chronic bronchitis, or chronic obstructive pulmonary disease (COPD) 5
- Pneumonia 6
- Infection 7
- Other condition 8 specify: _____
- Unknown 9

11. What was the date of the hospitalization:

/ /
Month Day Year

12. What was the name and location of the hospital?

13. Was (insert decedent's name) seen by a doctor at an emergency room or in any other facility in the last four weeks prior to death?

- No 0 Skip to question 15
- Yes 1
- Unknown 9 Skip to question 15

13a. What was the reason for this visit to an emergency room or doctors office? (select all that apply)

- Heart attack or heart disease 1
- Stroke 2
- Heart surgery 3
- Surgical procedure (other than heart) 4
- Emphysema, chronic bronchitis, or chronic obstructive pulmonary disease (COPD) 5
- Pneumonia 6
- Infection 7
- Other condition 8 specify: _____
- Unknown 9

14. What was the name and address of this physician or emergency room?

SYMPTOMS

“The next set of questions deals specifically with acute symptoms such as pain, discomfort that (insert decedent’s name) may have experienced at the time of his/her death.”

15. Did s/he experience pain, discomfort or tightness in the chest, left arm or jaw?

- No 0 Skip to question 22
Yes 1
Unknown 9 Skip to question 22

16. Did the pain, discomfort or tightness specifically involve the chest?

- No 0
Yes 1
Unknown 9

16a. Did (insert decedent’s name) ever take nitroglycerin for this pain?

- No 0
Yes 1
Unknown 9

17. Were these episodes new, or had they occurred previously?

- New symptoms 1 Skip to question 22
Previous symptoms 2
Unknown 9

18. Were the episodes getting longer or more frequent?

- No 0
Yes 1
Unknown 9

19. Were the episodes getting more severe?

- No 0
Yes 1
Unknown 9

****If No or Unknown to Questions 18 and 19, skip to Question 21****

20. Over what period of time did these episodes become longer, more frequent, or more severe?

- Days 1
Weeks 2
Months 3

Unknown 9

21. Did s/he experience shortness of breath?

No 0 Skip to item 22

Yes 1

Unknown 9 Skip to item 22

21a. Did s/he have shortness of breath while at rest?

No 0

Yes 1

Unknown 9

22. *“I apologize if this question sounds hard or if it makes you uncomfortable. Please be assured we respect your feelings about this unfortunate event. How long was it from (insert decedent’s name) last episode of symptoms to the time that s/he stopped breathing on his/her own?”*

Less than 5 minutes 1

Less than 1 hour 2

Less than 24 hours 3

Greater than 24 hours 4

Unknown 9

EMERGENCY MEDICAL CARE

“The next few questions are concerned with emergency medical care (insert decedent’s name) may have received prior to or at the time of death. You may have already given this information in an answer to an earlier question. Since it is important to obtain information specifically on emergency medical care, I hope you don’t mind if these questions seem repetitive.”

23. Was a physician, ambulance or other emergency medical team called?

No 0 Skip to question 24

Yes 1

Unknown 9 Skip to question 24

23a. How long was it from the time the last episode of symptoms started to the time that medical assistance was called for?

5 minutes or less 1

10 minutes or less 2

1 hour or less 3

6 hours or less 4

24 hours or less 5

More than 24 hours 6

Unknown 9

23b. How long was it from the time medical care was called to the time when it arrived?

5 minutes or less 1
10 minutes or less 2
1 hour or less 3
6 hours or less 4
24 hours or less 5
More than 24 hours 6
Unknown 9

24. Were resuscitation measures, such as CPR attempted?

No 0 Skip to question 25
Yes 1
Unknown 9 Skip to question 25

24a. Who started the CPR or resuscitation?

Bystander 1
Physician 2
Ambulance personnel 3
Fireman or Police 4
The informant 5
Other 6
Unknown 9

25. Was (insert decedent's name) taken to the hospital, emergency room or any other emergency care facility?

No 0
Yes 1
Unknown 9

26. Is there anyone else we could contact who might be able to provide additional information about the circumstances surrounding (insert decedent's name) death or his/her usual state of health?

No 0 Skip to Closing Script
Yes 1
Unknown 9 Skip to Closing Script

27. How is s/he related to the deceased?

- Spouse 1
- Daughter/Son 2
- Parent 3
- Friend 4
- Workmate 5
- Other relative 6 Specify: _____
- Other 7 Specify: _____

28. What is the name and address of this person?

CLOSING SCRIPT

“Thank you very much for your assistance in this study. Do you have any questions? Thanks again for your help.”

RELIABILITY

(To be completed after the interview)

29. On the basis of these questions, give your rating of reliability of the interview.

- Poor 1
- Fair 2
- Good 3



HCHS/SOL Physician Questionnaire

ID NUMBER:										
---------------	--	--	--	--	--	--	--	--	--	--

FORM CODE: PQE
VERSION: A
12/02/2008

Contact Occasion	0	1	SEQ #		
---------------------	---	---	-------	--	--

Administrative Information

0a. Completion Date: / /
month may year

0b. Staff ID:

Instructions: Please complete the following questions to the best of your ability by filling in the appropriate bubbles or writing the answer in the blank provided. Please return completed forms in the self addressed stamped envelope provided to the local HCHS/SOL field center.

Public reporting burden for this collection of information is estimated to average 30 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to, a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to: NIH, Project Clearance Branch, 6705 Rockledge Drive, MSC 7974, Bethesda, MD 20892-7974, ATTN: PRA (0925-0584). Do not return the completed form to this address.

DETAILS OF DEATH

1. Are you familiar with the events surrounding the decedent's death?

No 0
Yes 1

2. Did you witness the death?

No 0
Yes 1

If informant answered "Yes" to one or both of Items 1 and 2, please skip to Item 4.

3. If you answered "No" to both Questions, are you aware of another physician who could provide information regarding the death?

No 0 Please sign and date the bottom of this form
Yes 1

3a. Provide contact information. Please then sign and date the bottom of this form.

Name of physician: _____

Address: _____

CIRCUMSTANCES SURROUNDING DEATH

4. What do you believe to be the underlying cause of death?

- | | | |
|---|---|---|
| Acute Myocardial Infarction | 1 | <input type="checkbox"/> |
| Other Ischemic Heart Disease | 2 | <input type="checkbox"/> |
| Cerebrovascular Disease | 3 | <input type="checkbox"/> |
| Other Cardiovascular Disease | 4 | <input type="checkbox"/> |
| Emphysema, chronic bronchitis or chronic obstructive pulmonary disease (COPD) | 5 | <input type="checkbox"/> |
| Pneumonia | 6 | <input type="checkbox"/> |
| Asthma | 7 | <input type="checkbox"/> |
| Other Lung Disease | 8 | <input type="checkbox"/> specify: _____ |
| Non Cardio - Pulmonary Disease | 9 | <input type="checkbox"/> specify: _____ |

5. Please specify the time between the onset of the acute episode of symptoms and death. (We are defining death as the point where spontaneous breathing ceased and the patient never recovered.) Please check the appropriate time period.

- | | | |
|---------------------|---|--------------------------|
| Less than 5 minutes | 1 | <input type="checkbox"/> |
| 5 minutes to 1 hour | 2 | <input type="checkbox"/> |
| 1 hour to 24 hours | 3 | <input type="checkbox"/> |
| More than 24 hours | 4 | <input type="checkbox"/> |
| 1 day to 3 days | 5 | <input type="checkbox"/> |
| More than 3 days | 6 | <input type="checkbox"/> |
| Unknown | 9 | <input type="checkbox"/> |

6. Was there an acute episode of pain in the chest, left arm or jaw during the last 72 hours prior to death?

- | | | |
|---------|---|--------------------------|
| No | 0 | <input type="checkbox"/> |
| Yes | 1 | <input type="checkbox"/> |
| Unknown | 9 | <input type="checkbox"/> |

7. Was there an acute episode of shortness of breath during the 72 hours prior to death?

- | | | |
|---------|---|--------------------------|
| No | 0 | <input type="checkbox"/> |
| Yes | 1 | <input type="checkbox"/> |
| Unknown | 9 | <input type="checkbox"/> |

8. Was there an acute episode of wheezing during the 72 hours prior to death?

- | | | |
|---------|---|--------------------------|
| No | 0 | <input type="checkbox"/> |
| Yes | 1 | <input type="checkbox"/> |
| Unknown | 9 | <input type="checkbox"/> |

9. Did the decedent take or was s/he given nitrates or nitroglycerin at the time of the acute episode?

No 0
Yes 1
Unknown 9

MEDICAL HISTORY

10. Are you familiar with the decedent's medical history?

No 0 End questionnaire
Yes 1

11. Did the decedent have a medical history of any of the following conditions prior to the acute event which led to death?

11a. Myocardial Infarction (MI)?

No 0 Skip to 11b
Yes 1
Unknown 9 Skip to 11b

i. Date of most recent MI: //
month day year

11b. Angina Pectoris, Coronary Insufficiency or Other Chronic Ischemic Heart Disease?

No 0 Skip to 11c
Yes 1
Unknown 9 Skip to 11c

i. Date of first diagnosis: //
month day year

11c. Congestive Heart Failure (CHF) or Congestive Cardiomyopathy?

No 0 Skip to 11d
Yes 1
Unknown 9 Skip to 11d

i. Date of first exacerbation: //
month day year

11d. Stroke (CVA)?

No 0 Skip to 11e
Yes 1
Unknown 9 Skip to 11e

i. Date of most recent CVA: / /
month day year

11e. Transient Ischemic Attack (TIA)?

No 0 Skip to 11f
Yes 1
Unknown 9 Skip to 11f

i. Date of first diagnosis: / /
month day year

11f. Intermittent Claudication or Other Peripheral Arterial Disease (PAD)?

No 0 Skip to 11g
Yes 1
Unknown 9 Skip to 11g

11g. Lower Extremity Bypass, Angioplasty or Amputation Secondary to PAD?

No 0 Skip to 11h
Yes 1
Unknown 9 Skip to 11h

11h. Coronary Bypass Surgery?

No 0
Yes 1
Unknown 9

11i. Coronary Angioplasty?

No 0
Yes 1
Unknown 9

11j. Emphysema, chronic bronchitis, or Chronic Obstruction Pulmonary Disease (COPD)?

No 0 Skip to 11k
Yes 1
Unknown 9 Skip to 11k

i. Date of first exacerbation (or onset): / /



HCHS/SOL DEATH CERTIFICATE FORM

ID NUMBER:	<input type="text"/>	FORM CODE: DTH	Contact	<input type="text"/>	SEQ #	<input type="text"/>
	<input type="text"/>	VERSION: A 05/26/10	Occasion	<input type="text"/>		<input type="text"/>

Administrative Information

0a. Completion Date: / / Month Day Year

0b. Staff ID:

Event ID:

Event Date: / / Month Day Year

Instructions: The Death Certificate Form is completed for each death reported from the Annual Follow-up Form. A Death Certificate must be requested and obtained prior to completing this form.

1. Was a death certificate obtained? No 0 *skip out of form* Yes 1

2. Date of death: / / Month Day Year

3. Time of death: : 3a. 1 = A.M., 2 = P.M.

4. Did the decedent die in a hospital? No 0 *Skip to 6* Yes 1 Unknown 9

5. Was the death classified as: *(select one)*

- 1. dead on arrival (DOA)
- 2. emergency dept (ED)
- 3. outpatient
- 4. inpatient
- 5. none of the above
- 6. not recorded

6. Was this a coroner's or medical examiner's case? No 0 *Skip to 10* Yes 1

7. Was the name and address of the Coroner or medical examiner recorded? No 0 *Skip to 10* Yes 1

8. Name: _____

9. Address: a. Street _____

b. City _____ State _____ Zip code _____

c. Country _____

10. Was an autopsy performed? No 0 Yes 1

11. ICD-10 Code for **UNDERLYING** cause of death: .

12. All listed **ICD-10 Codes** for death:

a. .

e. .

h. .

b. .

f. .

i. .

c. .

g. .

j. .

d. .

13. Are there causes of death recorded on the death certificate? No 0 *skip to 14* Yes 1

13a. Immediate cause: _____

13b. Due to or as a consequence of (1) _____

13c. Due to or as a consequence of (2) _____

13d. Due to or as a consequence of (3) _____

14. Are there other significant conditions Recorded on the death certificate? No 0 *Skip to 16* Yes 1

15. Conditions: _____

16. Interval between onset and death for immediate cause of death:

- 1 = 5 minutes or less
- 2 = 1 hour or less
- 3 = 1 day or less
- 4 = 1 week or less

- 5 = 1 month or less
- 6 = more than 1 month
- 7 = unknown or not recorded

17. Was the name and address of the informant recorded? No 0 *Skip to 22* Yes 1

18. Name: _____

19. Address: a. Street _____

b. City _____ State _____ Zip code _____

c. Country _____

20. Relationship of informant to deceased: 1 = spouse, 2 = other, 3 = unknown

21. If other, specify: _____

22. Was the name and address of the certifying physician recorded? No 0 Yes 1

23. Name: _____

24. Address: a. Street _____

b. City _____ State _____ Zip code _____

c. Country _____

APPENDIX D Abstraction Forms

Myocardial Infarction Abstraction Form (MIF)

Heart Failure Abstraction Form (HTF)

Pulmonary Abstraction Form (PUL)



HCHS/SOL MYOCARDIAL INFARCTION ABSTRACTION FORM (MIF)

ID NUMBER:	<input type="text"/>	FORM CODE: MIF	Contact Occasion	<input type="text"/>	SEQ #	<input type="text"/>
	<input type="text"/>	VERSION: A 08/04/11		<input type="text"/>		<input type="text"/>

ADMINISTRATIVE INFORMATION

0A. Completion Date: / /
Month/Day/Year

0B. Staff ID:

Event ID:

Event Date: / /

Instructions: Answers are derived from the medical records received. Do not complete this form until all records are received (or classified as unobtainable) as indicated on the Verification of ICD Discharge Codes Form

A. GENERAL INFORMATION

1. Was the event (choose one):
 1= In hospital only 2= Emergency Dept. visit only(ED) 3= Both ED and in hospital

2. Date of arrival: (mm/dd/yyyy)

a. Time of arrival : *1= A.M., 2 = P.M.*

b. Date of admission

3. Date of discharge: (mm/dd/yyyy)

a. Time of discharge : *1= A.M., 2 = P.M.*

4. What was the primary admitting diagnosis code?

5. What was the primary discharge diagnosis code?

6. Did an emergency medical service unit transport the patient to this hospital? No/NR Yes

7. Was the patient transferred to this hospital from another hospital? No Yes

8. Was the patient's code status ever "no-code" or "DNR" (do not resuscitate)? No Yes

9. Was the patient alive at discharge? *If Yes, go to Item 10* No Yes

9.a. Was the patient dead on arrival? No Yes

9.b. Did the patient die in the Emergency Department? No Yes

9.c. Was an autopsy performed? No Yes

B. PRESENTING SIGNS AND SYMPTOMS

	<u>No</u>	<u>Yes</u>	<u>NR</u> Not recorded
10. Did the onset of the acute episode occur prior to admission?	0 <input type="checkbox"/>	1 <input type="checkbox"/>	9 <input type="checkbox"/>
a. If YES, estimate the time from onset of symptoms of acute condition to arrival at the hospital			Unsure <input type="checkbox"/>
< 1hr <input type="checkbox"/>		≥ 1- < 3 hrs <input type="checkbox"/>	
≥ 6 - < 12 hrs <input type="checkbox"/>		≥ 12 - < 24 hrs <input type="checkbox"/>	
		≥ 3 - < 6 hrs <input type="checkbox"/>	
		≥ 24 hrs. <input type="checkbox"/>	
11. Was there mention of an acute CHD event with onset <u>after</u> arrival at the hospital?	0 <input type="checkbox"/>	1 <input type="checkbox"/>	9 <input type="checkbox"/>
12. Was there an acute episode(s) of pain or discomfort (eg: tightness) anywhere in the chest, arm, shoulder throat or jaw, either within 72 hours prior to arrival to the hospital, or in conjunction with the in-hospital CHD event? (If No or NR, go to Item 13)	0 <input type="checkbox"/>	1 <input type="checkbox"/>	9 <input type="checkbox"/>
a. Did this pain or discomfort specifically involve the chest?	0 <input type="checkbox"/>	1 <input type="checkbox"/>	9 <input type="checkbox"/>
b. Did the pain get worse (crescendo) over time?	0 <input type="checkbox"/>	1 <input type="checkbox"/>	9 <input type="checkbox"/>
c. Was the pain or discomfort diagnosed as having a non-cardiac origin?	0 <input type="checkbox"/>	1 <input type="checkbox"/>	9 <input type="checkbox"/>
13. Was there nausea or vomiting associated with this event?	0 <input type="checkbox"/>	1 <input type="checkbox"/>	9 <input type="checkbox"/>
14. Was there diaphoresis associated with this event?	0 <input type="checkbox"/>	1 <input type="checkbox"/>	9 <input type="checkbox"/>
15. Was there fatigue or malaise associated with this event?	0 <input type="checkbox"/>	1 <input type="checkbox"/>	9 <input type="checkbox"/>
16. Vital Signs at arrival (or event onset) and not during CPR			
a. Blood pressure <input type="text"/> <input type="text"/> <input type="text"/> / <input type="text"/> <input type="text"/> <input type="text"/> mmHg			
b. Heart rate <input type="text"/> <input type="text"/> <input type="text"/> bpm			

C. MEDICAL HISTORY

	<u>No/NR</u>	<u>Yes</u>
17. Prior to this event was there history of any of the following:		
17.a. Myocardial infarction <i>If No or NR, skip to 17.b.</i>	0 <input type="checkbox"/>	1 <input type="checkbox"/>
1. If history of MI, then MI within 4 weeks of this event?	0 <input type="checkbox"/>	1 <input type="checkbox"/>
17.b. Angina	0 <input type="checkbox"/>	1 <input type="checkbox"/>
17.c. Percutaneous coronary intervention (PCI)	0 <input type="checkbox"/>	1 <input type="checkbox"/>
17.d. CABG	0 <input type="checkbox"/>	1 <input type="checkbox"/>
17.e. Coronary artery disease (CAD)	0 <input type="checkbox"/>	1 <input type="checkbox"/>
17.f. Heart failure	0 <input type="checkbox"/>	1 <input type="checkbox"/>
17.g. Arrhythmia	0 <input type="checkbox"/>	1 <input type="checkbox"/>

IF YES, specify type of arrhythmia

- | | | | |
|--------|--------------------------------------|----------------------------|----------------------------|
| 17.g.1 | Atrial Fibrillation/Flutter | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 17.g.2 | Ventricular Fibrillation/Tachycardia | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 17.g.3 | Other arrhythmia | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |

D. ACTIVE OR CURRENT MEDICAL PROBLEMS (DURING THIS HOSPITALIZATION)

18. Did a physician indicate any of these as being present during the hospitalization? Exclude old episodes; include only current conditions.

- | | <u>No/NR</u> | <u>Yes</u> |
|--|----------------------------|----------------------------|
| 18.a. Angina | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 18.b. Acute myocardial Infarction | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 18.c. ST elevation > 1mm with pain that is not present on ECG without pain | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 18.d. Congestive heart failure exacerbation or pulmonary edema | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 1. <u>IF YES</u> , Did heart failure/pulmonary edema occur within 24 hours of event onset? | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 18.e. Shock or cardiogenic shock | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 1. <u>IF YES</u> , Did shock occur within 24 hours of event onset? | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 18.f. Ventricular fibrillation, cardiac arrest or asystole | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 1. <u>IF YES</u> , Did the arrest occur within 24 hours of event onset? | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 18.g. Ventricular Tachycardia | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 18.h. Atrial fibrillation or atrial flutter | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |

E. BIOMARKERS

- | | <u>No/NR</u> | <u>Yes</u> |
|--|----------------------------|----------------------------|
| 19. Were cardiac enzymes reported within days 1-4 after arrival at the hospital or after the in-hospital CHD event?
If No/NR skip to 32 | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |

Biomarker Laboratory Standards:

***Units:** 1 = ng/mL 2 = Units/L 3 = µg/L

20. Range Set 1

- | | Upper limit of normal
(only) | Units* | N/A |
|-------------------|---|-------------------------|-------------------------|
| a. Total CK (CPK) | a. <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> . <input type="text"/> <input type="text"/> | b. <input type="text"/> | c. <input type="text"/> |
| b. CK-MB | a. <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> . <input type="text"/> <input type="text"/> | b. <input type="text"/> | c. <input type="text"/> |
| c. Total LDH | <input type="text"/> < a. <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> . <input type="text"/> <input type="text"/> | b. <input type="text"/> | c. <input type="text"/> |

f. Troponin < 1. 2. 3.

f.1. What type of Troponin was this?

a. Troponin, type not specified

b. Troponin I

c. Troponin T

d. High Sensitivity Troponin (HS)

e. Unsure

23. Day 1/Set 2 Date: / /

a. Total CK (CPK) 1. 2. 3.

b. CK-MB 1. 2. 3.

c. Total LDH 1. 2. 3.

d. LDH-1 1. 2. 3.

e. LDH-2 1. 2. 3.

f. Troponin < 1. 2. 3.

f.1. What type of Troponin was this?

a. Troponin, type not specified

b. Troponin I

c. Troponin T

d. High Sensitivity Troponin (HS)

e. Unsure

24. Day 2/Set 1 Date: / /

a. Total CK (CPK) 1. 2. 3.

b. CK-MB 1. 2. 3.

c. Total LDH 1. 2. 3.

d. LDH-1 1. 2. 3.

e. LDH-2 1. 2. 3.

f. Troponin < 1. 2. 3.

f.1. What type of Troponin was this?

a. Troponin, type not specified

b. Troponin I

c. Troponin T

d. High Sensitivity Troponin (HS)

e. Unsure

25. Day 2/Set 2 Date: / /

a. Total CK (CPK) 1. 2. 3.

b. CK-MB 1. 2. 3.

c. Total LDH 1. 2. 3.

d. LDH-1 1. 2. 3.

e. LDH-2 1. 2. 3.

f. Troponin < 1. 2. 3.

f.1. What type of Troponin was this?

a. Troponin, type not specified

b. Troponin I

c. Troponin T

d. High Sensitivity Troponin (HS)

e. Unsure

26. Day 3/Set 1 Date: / /

a. Total CK (CPK) 1. 2. 3.

b. CK-MB 1. 2. 3.

c. Total LDH 1. 2. 3.

d. LDH-1 1. 2. 3.

e. LDH-2 1. 2. 3.

f. Troponin < 1. 2. 3.

f.1. What type of Troponin was this?

a. Troponin, type not specified

b. Troponin I

c. Troponin T

d. High Sensitivity Troponin (HS)

e. Unsure

27. Day 3/Set 2 Date: / /

a. Total CK (CPK) 1. 2. 3.

b. CK-MB 1. 2. 3.

c. Total LDH 1. 2. 3.

d. LDH-1 1. 2. 3.

e. LDH-2 1. 2. 3.

f. Troponin < 1. 2. 3.

f.1. What type of Troponin was this?

a. Troponin, type not specified

b. Troponin I

c. Troponin T

d. High Sensitivity Troponin (HS)

e. Unsure

28. Day 4/Set 1 Date: / /

a. Total CK (CPK) 1. 2. 3.

b. CK-MB 1. 2. 3.

c. Total LDH 1. 2. 3.

d. LDH-1 1. 2. 3.

e. LDH-2 1. 2. 3.

f. Troponin < 1. 2. 3.

f.1. What type of Troponin was this?

a. Troponin, type not specified

b. Troponin I

c. Troponin T

d. High Sensitivity Troponin (HS)

e. Unsure

29. Day 4/Set 2 Date: //

Units* (see pg. 3) Range Set* (1 or 2) Words Code*

a. Total CK (CPK) 1. 2. 3.

b. CK-MB 1. 2. 3.

c. Total LDH 1. 2. 3.

d. LDH-1 1. 2. 3.

e. LDH-2 1. 2. 3.

f. Troponin < 1. 2. 3.

f.1. What type of Troponin was this?

a. Troponin, type not specified

b. Troponin I

c. Troponin T

d. High Sensitivity Troponin (HS)

e. Unsure

30. Is there mention of the patient having either trauma, a surgical procedure, or rhabdomyolysis, within one week prior to measurement of biomarkers? 0 No/NR 1 Yes
 If No/NR skip to 31

If yes, Indicate the type of procedure or trauma:

	No	Yes	Date	
a. Cardiac procedure	0 <input type="checkbox"/>	1 <input type="checkbox"/>	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> / <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>	c1. Specify: _____ f1. Specify: _____
b. CPR or cardioversion	0 <input type="checkbox"/>	1 <input type="checkbox"/>	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> / <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>	
c. Other cardiac trauma	0 <input type="checkbox"/>	1 <input type="checkbox"/>	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> / <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>	
d. Rhabdomyolysis	0 <input type="checkbox"/>	1 <input type="checkbox"/>	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> / <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>	
e. Intramuscular Injection	0 <input type="checkbox"/>	1 <input type="checkbox"/>	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> / <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>	
f. Non-cardiac procedure	0 <input type="checkbox"/>	1 <input type="checkbox"/>	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> / <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>	
g. Non-cardiac trauma	0 <input type="checkbox"/>	1 <input type="checkbox"/>	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> / <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>	

31. Enter the item number from the biomarkers section (items 22—30) of this form which corresponds to the first biomarker measurement performed after the trauma, cardiac procedure or rhabdomyolysis:

32. Is there evidence of hemolytic disease during the hospitalization? 0 No/NR 1 Yes

33. Did the participant have any active liver disease (cirrhosis, hepatitis, liver cancer, etc.)? 0 No/NR 1 Yes

F. Electrocardiography

34. Were any 12 lead ECGs taken during this admission? 0 No 1 Yes 9 NR
 (If this is an in-hospital event, then first ECG is at time of event)

- a. First ECG Date: / b. Copy of ECG enclosed? No 0 Yes 1
- c. Second ECG Date: / d. Copy of ECG enclosed? No 0 Yes 1
- e. Third ECG Date: / f. Copy of ECG enclosed? No 0 Yes 1
- g. Last ECG Date: / h. Copy of ECG enclosed? No 0 Yes 1

G. Procedures and Diagnostics

Were any of the following special procedures or operations performed during this hospitalization?
 (Mark all that apply)

- | | <u>No/NR</u> | <u>Yes</u> |
|---|----------------------------|----------------------------|
| 35. Transthoracic echocardiogram (TTE) performed? If No/NR, skip to 36 | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| a. LV Ejection fraction: <input type="text"/> <input type="text"/> % | | |
| 36. Was a Nuclear Medicare Scan (MUGA, SPECT or radionuclide ventriculogram (RVG)) performed? If No/NR, skip to 37 | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| a. Ejection fraction: LV: <input type="text"/> <input type="text"/> % b. RV: <input type="text"/> <input type="text"/> % | | |
| c. Stress test positive for ischemia | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 37. Was any stress test (treadmill, pharmacologic, or nuclear medicine) performed during this admission: If No/NR, skip to 38 | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| a. Ejection fraction: LV: <input type="text"/> <input type="text"/> % | | |
| b. Stress test positive for ischemia) | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| c. Greater than or equal to 1mm ST depression or elevation | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| d. Ischemic pain or equivalent occurred | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 38. Was a coronary angiography performed? If No/NR, skip to 37 | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| a. Date: (mm/dd/yyyy) <input type="text"/> / <input type="text"/> / <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> | | |
| b. Ejection fraction: LV: <input type="text"/> <input type="text"/> % | | |

No

Yes NR

c. 70% or greater obstruction of any coronary artery 0 1 9

d. Were coronary bypass grafts present? 0 1 9

1. If yes, number of occluded grafts:

H. Treatment

No/NR Yes

39. Was coronary reperfusion (CABG, PCI, thrombolysis) attempted? 0 1
If No/NR, Skip to 40

39.a. If yes, what was the approximate time from event onset to reperfusion?

< 2 hours 2-4 hours 4-6 hours 6-12 hours 12-24 hours > 24 hours not sure

40. Where any of the following treatments given during this hospitalization?

a. Coronary artery bypass graft surgery (CABG) 0 1

1. If yes, Date: / b. Time : c. 1= am, 2 = pm

b. Coronary atherectomy 0 1

1. If yes, Date: / b. Time : c. 1= am, 2 = pm

c. Intra-arterial or intravenous thrombolytic 0 1

1. If yes, Date: / b. Time : c. 1= am, 2 = pm

d. Coronary angioplasty without stent 0 1

1. If yes, Date: / b. Time : c. 1= am, 2 = pm

e. Coronary angioplasty with stent placement 0 1

1. If yes, Date: / b. Time : c. 1= am, 2 = pm

No/NR Yes

f. Valve surgery 0 1

g. Non-cardiac surgery 0 1

h. Aortic balloon pump 0 1

i. Pacemaker placement (temporary or permanent) 0 1

No/NR Yes

j. Cardioversion or defibrillation

0

1

1. If yes, Date: /

b. Time :

c. 1= am, 2 = pm

2. If cardioversion took place after arrival at the hospital, what rhythm(s) were present prior to cardioversion?

	<u>No/NR</u>	<u>Yes</u>
a. Ventricular Fibrillation/Flutter	0 <input type="checkbox"/>	1 <input type="checkbox"/>
b. Ventricular Tachycardia (VT)	0 <input type="checkbox"/>	1 <input type="checkbox"/>
c. Asystole	0 <input type="checkbox"/>	1 <input type="checkbox"/>
d. Complete AV Block (3 HB)	0 <input type="checkbox"/>	1 <input type="checkbox"/>
e. Atrial Fibrillation/Flutter	0 <input type="checkbox"/>	1 <input type="checkbox"/>
f. Pulseless Electrical Activity (PEA)	0 <input type="checkbox"/>	1 <input type="checkbox"/>

41. During the hospitalization or at discharge, did the participant receive any of the following medications?

	Admission Meds		Discharge Meds	
	<u>No/NR</u>	<u>Yes</u>	<u>No/NR</u>	<u>Yes</u>
a. Nitroglycerin	0 <input type="checkbox"/>	1 <input type="checkbox"/>	0 <input type="checkbox"/>	1 <input type="checkbox"/>
b. Beta Blockers	0 <input type="checkbox"/>	1 <input type="checkbox"/>	0 <input type="checkbox"/>	1 <input type="checkbox"/>
c. Calcium Channel Blockers	0 <input type="checkbox"/>	1 <input type="checkbox"/>	0 <input type="checkbox"/>	1 <input type="checkbox"/>
d. ACE Inhibitor or ARB	0 <input type="checkbox"/>	1 <input type="checkbox"/>	0 <input type="checkbox"/>	1 <input type="checkbox"/>
d. Scheduled aspirin (not PRN)	0 <input type="checkbox"/>	1 <input type="checkbox"/>	0 <input type="checkbox"/>	1 <input type="checkbox"/>
e. Heparin or Enoxaparin	0 <input type="checkbox"/>	1 <input type="checkbox"/>	0 <input type="checkbox"/>	1 <input type="checkbox"/>
f. Coumadin, warafin, panwafarin, dicumarol	0 <input type="checkbox"/>	1 <input type="checkbox"/>	0 <input type="checkbox"/>	1 <input type="checkbox"/>
g. Anti-platelet agents (non-aspirin)	0 <input type="checkbox"/>	1 <input type="checkbox"/>	0 <input type="checkbox"/>	1 <input type="checkbox"/>
h. Statin	0 <input type="checkbox"/>	1 <input type="checkbox"/>	0 <input type="checkbox"/>	1 <input type="checkbox"/>

42. During this hospitalization was this patient treated with:

	<u>No/NR</u>	<u>Yes</u>
a. IV pressors	0 <input type="checkbox"/>	1 <input type="checkbox"/>
b. IV nitroglycerin	0 <input type="checkbox"/>	1 <input type="checkbox"/>
c. IIb / IIIa inhibitors or thrombin inhibitors	0 <input type="checkbox"/>	1 <input type="checkbox"/>



HCHS/SOL HEART FAILURE ABSTRACTION FORM (HTF)

PARTICIPANT ID NUMBER:	<input style="width: 20px; height: 20px;" type="text"/>
---------------------------	---

FORM CODE:
HTF
VERSION: A
7/26/11

<input style="width: 20px; height: 20px;" type="text"/> <input style="width: 20px; height: 20px;" type="text"/>	<input style="width: 20px; height: 20px;" type="text"/> <input style="width: 20px; height: 20px;" type="text"/>
---	---

ADMINISTRATIVE INFORMATION

0B. Completion Date: / /
Month Day Year

0B. Staff ID:

Event ID:

Event Date: / /

Instructions: Answers are derived from the medical records received. Do not complete this form until all records are received (or classified as unobtainable) as indicated on the Verification of ICD Discharge Codes Form

A. GENERAL INFORMATION

1. Was the event (choose one):
 1= In hospital
 2= Emergency Dept. (ED)
 3= Both ED and in hospital

2. Date of arrival: (mm/dd/yyyy) / /

a. Time of arrival : 1 = A.M., 2 = P.M.

b. Date of admission / /

3. Date of discharge: (mm/dd/yyyy) / /

a. Time of discharge : 1 = A.M., 2 = P.M.

4. What was the primary admitting diagnosis code? .

5. What was the primary discharge diagnosis code? .

6. Did an emergency medical service unit transport the patient to this hospital?
No/NR

Yes

7. Was the patient transferred to this hospital from another hospital? 0 1
8. Was the patient's code status ever "no-code" or "DNR" (do not resuscitate)? 0 1
9. Was the patient alive at discharge? 0 1

B. SIGNS AND SYMPTOMS

I. Signs and Symptoms

10. Did the patient have any of the following signs or symptoms at the time of event ?

	<u>No</u>	<u>Yes</u>	<u>NR</u> Not
recorded			
a. An increase or new onset of paroxysmal nocturnal dyspnea (PND)	0 <input type="checkbox"/>	1 <input type="checkbox"/>	9 <input type="checkbox"/>
b. An increase or new onset of orthopnea?	0 <input type="checkbox"/>	1 <input type="checkbox"/>	9 <input type="checkbox"/>
c. An increase or new onset of shortness of breath?	0 <input type="checkbox"/>	1 <input type="checkbox"/>	9 <input type="checkbox"/>
d. An increase or new onset edema?	0 <input type="checkbox"/>	1 <input type="checkbox"/>	9 <input type="checkbox"/>
e. Hypoxia	0 <input type="checkbox"/>	1 <input type="checkbox"/>	9 <input type="checkbox"/>
f. Dyspnea (at rest) or tachypnea (RR>22)	0 <input type="checkbox"/>	1 <input type="checkbox"/>	9 <input type="checkbox"/>
g. Dyspnea (walking or on exertion)	0 <input type="checkbox"/>	1 <input type="checkbox"/>	9 <input type="checkbox"/>

II. Evidence in Physicians' Notes of Reason for Event

11. Was there evidence in the doctor's notes that the reason for this event was an exacerbation of heart failure?	0 <input type="checkbox"/>	1 <input type="checkbox"/>	9 <input type="checkbox"/>
12. Did the patient have new onset or progressive signs/symptoms of heart failure prior to presentation in ED or hospital?	0 <input type="checkbox"/>	1 <input type="checkbox"/>	9 <input type="checkbox"/>
13. Did the physician's note or discharge summary indicate the presence of any of the following specific types of heart failure? (check all that apply)	<u>No</u>	<u>Yes</u>	
a. Diastolic heart failure	0 <input type="checkbox"/>	1 <input type="checkbox"/>	
b. Systolic heart failure	0 <input type="checkbox"/>	1 <input type="checkbox"/>	
c. Right-sided heart failure	0 <input type="checkbox"/>	1 <input type="checkbox"/>	
d. Ischemic cardiomyopathy	0 <input type="checkbox"/>	1 <input type="checkbox"/>	
e. Idiopathic/dilated cardiomyopathy	0 <input type="checkbox"/>	1 <input type="checkbox"/>	
f. Myocarditis	0 <input type="checkbox"/>	1 <input type="checkbox"/>	

- g. Peripartum cardiomyopathy 0 1
- h. Other specific cardiomyopathy/heart failure 0 1

1. If other cardiomyopathy, specify type _____

III. Prior cardiac testing

14. Was cardiac imaging performed prior to this hospitalization?
 No/NR 0 Yes 1

skip to 15

a. Lowest LV ejection fraction recorded: %

b. Qualitative description of ejection fraction:

- Normal..... N
- Mildly reduced.....M
- Decreased moderately.....D
- Severely reduced.....S
- None of the above.....O
- Unsure-Not available.....U

c. Time (months) since recording of lowest ejection fraction: . months

d. Type of Imaging from which ejection fraction was obtained:

- 1. ECHO
- 2. MUGA
- 3. Catheterization with ventriculography
- 4. CT
- 5. MRI
- 6. Other
- 7. Unknown

C. MEDICAL HISTORY

	<u>No/NR</u>	<u>Yes</u>
15. Prior to this event was there a history of any of the following:		
a. Diagnosis of heart failure	0 <input type="checkbox"/>	1 <input type="checkbox"/>
b. Prior hospitalization for heart failure	0 <input type="checkbox"/>	1 <input type="checkbox"/>
c. Treatment for heart failure	0 <input type="checkbox"/>	1 <input type="checkbox"/>
d. Valvular heart disease	0 <input type="checkbox"/>	1 <input type="checkbox"/>
e. Rheumatic heart disease (RHD)	0 <input type="checkbox"/>	1 <input type="checkbox"/>
f. Congenital heart disease	0 <input type="checkbox"/>	1 <input type="checkbox"/>
g. Coronary heart disease (ever)	0 <input type="checkbox"/>	1 <input type="checkbox"/>

h. Coronary heart disease (within year)	0 <input type="checkbox"/>	1 <input type="checkbox"/>
i. Angina	0 <input type="checkbox"/>	1 <input type="checkbox"/>
j. Myocardial infarction	0 <input type="checkbox"/>	1 <input type="checkbox"/>
	<u>No/NR</u>	<u>Yes</u>
k. Atrial fibrillation/atrial flutter	0 <input type="checkbox"/>	1 <input type="checkbox"/>
l. Heart block or other bradycardia	0 <input type="checkbox"/>	1 <input type="checkbox"/>
m. Ventricular fibrillation or tachycardia	0 <input type="checkbox"/>	1 <input type="checkbox"/>
n. Hypertension	0 <input type="checkbox"/>	1 <input type="checkbox"/>
o. Diabetes	0 <input type="checkbox"/>	1 <input type="checkbox"/>
p. Chronic Obstructive Pulmonary Disease (COPD)	0 <input type="checkbox"/>	1 <input type="checkbox"/>
q. Cor pulmonale	0 <input type="checkbox"/>	1 <input type="checkbox"/>
r. Pulmonary hypertension	0 <input type="checkbox"/>	1 <input type="checkbox"/>

D. SURGICAL HISTORY

16. Past cardiac procedures	<u>No/NR</u>	<u>Yes</u>
a. CABG	0 <input type="checkbox"/>	1 <input type="checkbox"/>
b. Percutaneous coronary intervention (PCI)	0 <input type="checkbox"/>	1 <input type="checkbox"/>
c. Valve surgery	0 <input type="checkbox"/>	1 <input type="checkbox"/>
d. Pacemaker	0 <input type="checkbox"/>	1 <input type="checkbox"/>
e. Automatic Internal Cardiac Defibrillator (AICD)	0 <input type="checkbox"/>	1 <input type="checkbox"/>
f. Ablation for arrhythmia	0 <input type="checkbox"/>	1 <input type="checkbox"/>
g. Cardiac transplant	0 <input type="checkbox"/>	1 <input type="checkbox"/>
h. Ventricular Assist Device (VAD)	0 <input type="checkbox"/>	1 <input type="checkbox"/>

E. HOSPITAL COURSE

17. Current or Active Problems	<u>No/NR</u>	<u>Yes</u>
--------------------------------	--------------	------------

- | | | |
|---|----------------------------|----------------------------|
| a. Myocardial Infarction | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| b. Shock or Cardiogenic Shock | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| c. Ventricular Fibrillation, Cardiac Arrest or Asystole | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| d. Ventricular Tachycardia | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| e. Atrial Fibrillation/Atrial Flutter | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| | <u>No/NR</u> | <u>Yes</u> |
| f. COPD exacerbation | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| g. Cardiac Surgery – CABG or valvular surgery | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| h. Non-cardiac surgery | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| i. Pulmonary Embolus | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| j. Pneumonia | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |

F. PHYSICAL EXAM

18. Vital Signs at Admission (or at onset of event)

- | | | |
|----------------------------------|--|--|
| a. First available weight or BMI | <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> | <input type="checkbox"/> 1= weight in lbs
2= weight in Kg
3= BMI |
|----------------------------------|--|--|

19. Did the patient have any of the following signs?

- | | <u>No</u> | <u>Yes</u> | <u>NR</u>
Not |
|---|----------------------------|----------------------------|----------------------------|
| recorded a. Jugular venous distension (JVD) | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> | 9 <input type="checkbox"/> |
| b. Heart/Lung sounds: | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> | 9 <input type="checkbox"/> |
| 1. crackles or rales | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> | 9 <input type="checkbox"/> |
| 2. wheezing | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> | 9 <input type="checkbox"/> |
| 3. rhonchi | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> | 9 <input type="checkbox"/> |
| 4. S3 gallop | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> | 9 <input type="checkbox"/> |
| c. Lower extremity edema-unilateral | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> | 9 <input type="checkbox"/> |
| d. Lower extremity edema-bilateral | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> | 9 <input type="checkbox"/> |

G. DIAGNOSTIC TESTS

20. Was a chest X-ray performed during this event? No/NR 0 *skip to 21* Yes 1

21. Did the patient have any of the following signs on chest x-ray at any time during this event?

	<u>No/NR</u>	<u>Yes</u>
a. Pulmonary edema or CHF	0 <input type="checkbox"/>	1 <input type="checkbox"/>
b. Cardiomegaly or Cardiothoracic ratio ≥ 0.5	0 <input type="checkbox"/>	1 <input type="checkbox"/>

	<u>No/NR</u>	<u>Yes</u>
c. Pulmonary vascular congestion or Interstitial edema	0 <input type="checkbox"/>	1 <input type="checkbox"/>
d. Bilateral or unilateral pleural effusion	0 <input type="checkbox"/>	1 <input type="checkbox"/>

22. Was a chest/lung CT scan or CT angiogram (CTA) performed during this hospitalization? 0 1

23. Did the patient have any of the following signs on CT scan at any time during this hospitalization?

a. Pulmonary edema or pulmonary vascular congestion	0 <input type="checkbox"/>	1 <input type="checkbox"/>
b. Cardiomegaly	0 <input type="checkbox"/>	1 <input type="checkbox"/>
c. Bilateral or unilateral pleural effusion	0 <input type="checkbox"/>	1 <input type="checkbox"/>
d. Enlarged superior or inferior vena cava	0 <input type="checkbox"/>	1 <input type="checkbox"/>
e. Enlarged Pulmonary arteries	0 <input type="checkbox"/>	1 <input type="checkbox"/>

24. Was a transthoracic echocardiogram (TTE) performed? 0 1
Skip to 27

First transthoracic echocardiogram performed after onset of event:

a. Date (mm/dd/yyyy) / /

b. Left Ventricular Ejection Fraction: %

c. Record the following if present on transthoracic echocardiogram:

	<u>None</u>	<u>Mild</u>	<u>Mod</u>	<u>Severe</u>	<u>NR</u>
1. Left ventricular hypertrophy (LVH)	0 <input type="checkbox"/>	3 <input type="checkbox"/>	4 <input type="checkbox"/>	5 <input type="checkbox"/>	9 <input type="checkbox"/>

- 2. Impaired LV systolic function 0 3 4 5 9
- 3. Impaired RV systolic function 0 3 4 5 9
- 4. Pulmonary hypertension 0 3 4 5 9
- 5. Valvular heart disease 0 3 4 5 9

- 6. Diastolic dysfunction No/NR Yes
0 1
- 7. Stress test positive for ischemia 0 1
- 8. Regional wall motion abnormalities 0 1

- 9. Dilated left ventricle No/NR Yes
0 1
- 10. Dilated right ventricle 0 1

25. Was a transesophageal echocardiogram (TEE) performed? 0 1
skip to 26

First transesophageal echocardiogram performed after onset of event:

a. Date (mm/dd/yyyy) / /

b. Ejection fraction: b.1. LV % b.2. RV %

c. Record the following if present on transesophageal echocardiogram:

- | | <u>None</u> | <u>Mild</u> | <u>Moderate</u> | <u>Severe</u> | <u>NR</u> |
|----------------------------------|----------------------------|----------------------------|----------------------------|----------------------------|----------------------------|
| 1. Impaired LV systolic function | 0 <input type="checkbox"/> | 3 <input type="checkbox"/> | 4 <input type="checkbox"/> | 5 <input type="checkbox"/> | 9 <input type="checkbox"/> |
| 2. Impaired RV systolic function | 0 <input type="checkbox"/> | 3 <input type="checkbox"/> | 4 <input type="checkbox"/> | 5 <input type="checkbox"/> | 9 <input type="checkbox"/> |

- | | <u>No/NR</u> | <u>Yes</u> |
|---------------------------------------|----------------------------|----------------------------|
| 3. Regional wall motion abnormalities | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 4. Dilated left ventricle | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 5. Dilated right ventricle | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 6. Valvular heart disease | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |

No/NR Yes

26. Was coronary angiography performed? 0 1
skip to 27
- a. Date: (mm/dd/yyyy) // b. LV Ejection fraction: %
No/NR Yes
- c. 70% or greater obstruction of any coronary artery 0 1
No/NR Yes
27. Was a cardiac multiple-gated acquisition scan (MUGA) or RVG performed? 0 1
skip to 28
- a. Ejection fraction: LV: % b. RV: %
No/NR Yes
- c. Dilated ventricle or impaired ventricular function 0 1
28. Was a cardiac Magnetic Resonance Imaging (MRI) performed? 0 1
skip to 29
- a. Ejection fraction: LV: % b. RV: %
29. Did any imaging/diagnostic test show:
- a. Ejection fraction: LV: %
No/NR Yes
- b. Stress test positive for ischemia? 0 1
- c. Dilated ventricle or impaired ventricular function 0 1
- d. Left ventricular diastolic dysfunction 0 1
- e. Ventricular Septal Defect (VSD) 0 1
- f. Atrial Septal Defect (ASD) 0 1
- g. Patent Ductus Arteriosus (PDA) 0 1
- h. Artificial heart valve 0 1
- i. Hypertrophic Obstructive Cardiomyopathy (HOCM) 0 1
- j. Valvular Heart Disease 0 1

H. LABORATORY TESTS

	<u>a.Worst*</u>	<u>b.Last</u>	<u>c. Upper Limit Normal</u>
30. BNP (pg/mL)	□□□□□.□	□□□□□.□	□□□□□.□
31. ProBNP (pg/mL)	□□□□□.□	□□□□□.□	□□□□□.□
32. Troponin □	□ < a. □□□□□.□□		c. □ < □□□□□.□

a. If troponin value available, then what type of Troponin was this?

- | | |
|-----------------------------------|---|
| 1. Troponin, type not specified | □ |
| 2. Troponin I | □ |
| 3. Troponin T | □ |
| 4. High Sensitivity Troponin (HS) | □ |
| 5. Unsure | □ |

	<u>a.Worst*</u>
33. Sodium (mEq/L)	□□□
34. Serum creatinine (mg/dL)	□□.□□
35. BUN (mg/dL)	□□□
36. Hemoglobin (g/dL)	□□.□
37. Hematocrit (%)	□□.□

****Worst = highest value with exception of hemoglobin, hematocrit and sodium. For these the worst is the lowest value***

I. TREATMENTS

38. Were any of the following treatments given during this visit?	<u>No/NR</u>	<u>Yes</u>
a. Cardioversion or Defibrillation	0 □	1 □
b. Aortic balloon pump	0 □	1 □
c. Percutaneous coronary intervention (PCI)	0 □	1 □
d. CPAP or BIPAP	0 □	1 □
e. Mechanical Ventilation	0 □	1 □
f. Thoracentesis (therapeutic or diagnostic)	0 □	1 □

g. Ventricular Assist Device (VAD)

0

1

h. Heart transplant

0

1

J. MEDICATIONS

Prior to hospitalization

At discharge

	<u>No/NR</u>	<u>Yes</u>		<u>No/NR</u>	<u>Yes</u>
39. ACE inhibitors	0 <input type="checkbox"/>	1 <input type="checkbox"/>		0 <input type="checkbox"/>	1 <input type="checkbox"/>
40. Angiotensin II receptor Blockers	0 <input type="checkbox"/>	1 <input type="checkbox"/>		0 <input type="checkbox"/>	1 <input type="checkbox"/>
41. Beta blockers	0 <input type="checkbox"/>	1 <input type="checkbox"/>		0 <input type="checkbox"/>	1 <input type="checkbox"/>
42. Digitalis	0 <input type="checkbox"/>	1 <input type="checkbox"/>		0 <input type="checkbox"/>	1 <input type="checkbox"/>
43. Diuretics	0 <input type="checkbox"/>	1 <input type="checkbox"/>		0 <input type="checkbox"/>	1 <input type="checkbox"/>
				<u>No/NR</u>	<u>Yes</u>
44. Aldosterone blocker	0 <input type="checkbox"/>	1 <input type="checkbox"/>		0 <input type="checkbox"/>	1 <input type="checkbox"/>
45. Lipid lowering agents	0 <input type="checkbox"/>	1 <input type="checkbox"/>		0 <input type="checkbox"/>	1 <input type="checkbox"/>
46. Nitrates	0 <input type="checkbox"/>	1 <input type="checkbox"/>		0 <input type="checkbox"/>	1 <input type="checkbox"/>
47. Hydralazine	0 <input type="checkbox"/>	1 <input type="checkbox"/>		0 <input type="checkbox"/>	1 <input type="checkbox"/>
48. IV drugs during this hospitalization?				<u>No/NR</u>	<u>Yes</u>
a. IV inotropes				0 <input type="checkbox"/>	1 <input type="checkbox"/>
b. IV diuretics				0 <input type="checkbox"/>	1 <input type="checkbox"/>

SIGNS AND SYMPTOMS

I. Signs and Symptoms

11. Did the patient have any of the following signs or symptoms at the time of the event?

<u>Recorded</u>	<u>No</u>	<u>Yes</u>	<u>Not</u>
a. New onset or increase in cough?	0 <input type="checkbox"/>	1 <input type="checkbox"/>	9 <input type="checkbox"/>
b. New onset or increase in sputum production?	0 <input type="checkbox"/>	1 <input type="checkbox"/>	9 <input type="checkbox"/>
c. New onset or increase in sputum purulence?	0 <input type="checkbox"/>	1 <input type="checkbox"/>	9 <input type="checkbox"/>
d. New onset or increase in wheezing?	0 <input type="checkbox"/>	1 <input type="checkbox"/>	9 <input type="checkbox"/>
e. New onset or increase in chest tightness or chest pain?	0 <input type="checkbox"/>	1 <input type="checkbox"/>	9 <input type="checkbox"/>
f. New onset or increase in leg edema (unilateral or bilateral)?	0 <input type="checkbox"/>	1 <input type="checkbox"/>	9 <input type="checkbox"/>
g. New onset or increase in use of rescue bronchodilator?	0 <input type="checkbox"/>	1 <input type="checkbox"/>	9 <input type="checkbox"/>
h. New onset or increase in dyspnea?	0 <input type="checkbox"/>	1 <input type="checkbox"/>	9 <input type="checkbox"/>
i. Dyspnea (at rest)?	0 <input type="checkbox"/>	1 <input type="checkbox"/>	9 <input type="checkbox"/>
j. Dyspnea (walking or on exertion)?	0 <input type="checkbox"/>	1 <input type="checkbox"/>	9 <input type="checkbox"/>
k. Woken up at night by shortness of breath?	0 <input type="checkbox"/>	1 <input type="checkbox"/>	9 <input type="checkbox"/>
l. Fever?	0 <input type="checkbox"/>	1 <input type="checkbox"/>	9 <input type="checkbox"/>
m. Delirium or altered mental status (AMS)?	0 <input type="checkbox"/>	1 <input type="checkbox"/>	9 <input type="checkbox"/>

II. Evidence in Physicians' Notes of Reason for Event

	<u>No/NR</u>	<u>Yes</u>
12. Was there evidence in the doctor's notes that the reason for this event may be an exacerbation of COPD, chronic bronchitis, or emphysema?	0 <input type="checkbox"/>	1 <input type="checkbox"/>
13. Was there evidence in the doctor's notes that the reason for this event may be an exacerbation of asthma?	0 <input type="checkbox"/>	1 <input type="checkbox"/>
14. Did the patient have new onset or progressive signs/symptoms of this exacerbation Prior to presentation in ED or hospital?	0 <input type="checkbox"/>	1 <input type="checkbox"/>

B. MEDICAL HISTORY

15. Prior to this event was there a history of any of the following:	<u>No/NR</u>	<u>Yes</u>
a. Asthma	0 <input type="checkbox"/>	1 <input type="checkbox"/>
b. Chronic bronchitis	0 <input type="checkbox"/>	1 <input type="checkbox"/>
c. Emphysema	0 <input type="checkbox"/>	1 <input type="checkbox"/>
d. Chronic obstructive pulmonary disease (COPD)	0 <input type="checkbox"/>	1 <input type="checkbox"/>
e. Pulmonary fibrosis	0 <input type="checkbox"/>	1 <input type="checkbox"/>
f. Sarcoidosis	0 <input type="checkbox"/>	1 <input type="checkbox"/>
g. Lung cancer	0 <input type="checkbox"/>	1 <input type="checkbox"/>

15. Prior to this event was there a history of any of the following:
- | | <u>No/NR</u> | <u>Yes</u> |
|---|----------------------------|----------------------------|
| h. Lung resection or lobectomy | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| i. Home oxygen (do not include CPAP) | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| j. Pulmonary embolus..... | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| k. Pulmonary hypertension..... | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| l. Cor pulmonale | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| m. Obstructive Sleep Apnea (OSA)..... | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| n. Coronary artery disease | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| o. Heart failure | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| p. Atrial fibrillation/atrial flutter | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| q. Diabetes | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| r. Pulmonary Tuberculosis | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| s. Bronchiectasis | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |

16. If prior PFT results were provided, what is percent predicted FEV1?

16.a. Pre-bronchodilator .%

16.b. Post-bronchodilator .%

17. What is FEV₁/FVC ratio? . units

1=proportion or decimal
2=percent (If percent, then assure not percent predicted for the ratio)

C. HOSPITAL COURSE

18. Current or Active Problems anytime during this visit
- | | <u>No/NR</u> | <u>Yes</u> |
|--|----------------------------|----------------------------|
| a. Upper Respiratory Infection (sinusitis, nasopharyngitis, pharyngitis,
epiglottitis, laryngitis, laryngotracheitis, acute bronchitis) | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| b. Pneumonia | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| c. Pulmonary embolus | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| d. Myocardial infarction | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| e. Heart failure exacerbation | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| f. Atrial fibrillation/atrial flutter | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| g. Supraventricular Tachycardia (SVT) or multifocal atrial tachycardia (MAT) | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| h. Cardiac Surgery – CABG or Valvular Surgery | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| i. Non-cardiac surgery | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |

D. PHYSICAL EXAM

19. Vital Signs at arrival to hospital or ED (or at onset of event if began after arrival)

a. heart rate

bpm

- b. respiration rate per minute
- c. Oxygen Saturation (SpO₂/pulse oximetry) .%
- c.1. Oxygen Sats on room air? No 0 Yes 1 *Skip to 19d* Unknown 9 *Skip to 19d*
- c.2. If not on room air, what level oxygen? . 1= Liters, 2=Percent
- d. Weight . 1= Lbs, 2=Kg

20. Did the patient have any of the following signs (at the time of the event)?

- | | <u>No</u> | <u>Yes</u> | <u>NR</u> |
|--|----------------------------|----------------------------|----------------------------|
| a. Use of accessory muscles | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> | 9 <input type="checkbox"/> |
| b. Cyanosis | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> | 9 <input type="checkbox"/> |
| c. Clubbing | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> | 9 <input type="checkbox"/> |
| d. Jugular venous distention (JVD) or distended neck veins | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> | 9 <input type="checkbox"/> |
| e. Crackles/rales | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> | 9 <input type="checkbox"/> |
| f. Wheezing or rhonchi | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> | 9 <input type="checkbox"/> |
| g. Decreased <u>unilateral</u> breath sounds | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> | 9 <input type="checkbox"/> |
| h. Decreased <u>bilateral</u> breath sounds | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> | 9 <input type="checkbox"/> |
| i. Prolonged expiratory time | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> | 9 <input type="checkbox"/> |
| j. Egophony | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> | 9 <input type="checkbox"/> |
| k. Lower extremity edema (unilateral or bilateral) | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> | 9 <input type="checkbox"/> |

E. DIAGNOSTIC TESTS

21. Was a chest X-ray performed during this event? No/NR 0 *skip to 23* Yes 1

22. Did the patient have any of the following signs on chest x-ray at any time during this event?

- | | <u>No/NR</u> | <u>Yes</u> |
|--|----------------------------|----------------------------|
| a. Hyperinflation | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| b. Flattened diaphragms | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| c. Consolidation or infiltrate..... | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| d. Scarring | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| e. Nodule(s) > 8mm..... | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| f. Mass(es) > 3cm..... | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| g. Pulmonary edema, pulmonary vascular congestion (alveolar, interstitial) | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| h. Bilateral pleural effusion..... | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| i. Unilateral pleural effusion..... | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| j. Emphysema | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| k. Cardiomegaly | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |

23. Was a chest/lung CT scan or CT angiogram (CTA) performed during this event?

No/NR 0 *skip to 25* Yes 1

24. Did the patient have any of the following signs on CT scan at any time during this event?

- | | <u>No/NR</u> | <u>Yes</u> |
|---|----------------------------|----------------------------|
| a. Emphysema | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| b. Nodule(s) > 8mm | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| c. Mass(es) > 3cm | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| d. Lymphadenopathy..... | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| e. Ground glass changes | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| f. Pneumonia | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| g. Fibrosis or honeycombing..... | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| h. Filling defect—vascular (PE)..... | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| i. Filling defect—mucus plug..... | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| j. Cysts or blebs | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| k. Atelectasis..... | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| l. Calcifications..... | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| m. Pulmonary embolus | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| n. Enlarged pulmonary artery..... | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| o. Bronchiectasis..... | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| p. Pulmonary edema or pulmonary vascular congestion | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| q. Cardiomegaly | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| r. Bilateral pleural effusion..... | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| s. Unilateral pleural effusion..... | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| t. Airway wall thickening..... | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |

25. Was spirometry (lung function testing) performed during this hospitalization?

No/NR 0 *skip to 26* Yes 1

- | | | | | |
|-----|-----------------------------|--|--|--|
| a1 | FEV ₁ | <input type="text"/> <input type="text"/> . <input type="text"/> <input type="text"/> L | a2. FEV ₁ Percent Predicted | <input type="text"/> <input type="text"/> <input type="text"/> % |
| b1 | FVC | <input type="text"/> <input type="text"/> . <input type="text"/> <input type="text"/> L | b2. FVC Percent Predicted | <input type="text"/> <input type="text"/> <input type="text"/> % |
| c1. | FEV ₁ /FVC ratio | <input type="text"/> <input type="text"/> <input type="text"/> . <input type="text"/> <input type="text"/> units | <input type="checkbox"/> 1=proportion or decimal
<input type="checkbox"/> 2=percent (If percent, then assure not percent predicted for the ratio) | |

BIOCHEMICAL TESTS

31. White Blood Cell Count .

a. First (at event)

32. Hemoglobin (g/dL) .

33. Hematocrit (%) .

34. Sodium (mEq/L)

35. Serum creatinine (mg/dL) .

36. BUN (mg/dL) .

37. Bicarbonate (total CO₂) .

a. First (at event)

b. Upper limit normal

38. BNP (pg/mL) a. . b. .

39. ProBNP (pg/mL) a. . b. .

40. Were Arterial Blood Gases (ABGs) obtained? No/NR 0 *Skip to 41* Yes 1

a. First blood gas (at time of event)

b. Last blood gas

pH 1. .

1. .

PaCO₂ 2. . mmHg

2. . mmHg

PaO₂ 3. . mmHg

3. . mmHg

O₂ Saturation 4. . %

4. . %

c. Blood gas on room air? No 0 Yes 1 *Skip to 41* Not Recorded 9 *Skip to 41*

c.1. If not on room air, what level oxygen? . 1= Liters, 2=Percent

41. Was a sputum culture done? No/NR 0 *Skip to 42* Yes 1

a. Culture Results Neg 0 *Skip to 42* Pos 1 Not Recorded 9 *Skip to 42*

- b. If yes, were any of the following reported in the sputum culture?
- | | <u>No</u> | <u>Yes</u> |
|---|----------------------------|----------------------------|
| 1. Haemophilus Influenzae..... | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 2. Moraxella Catarrhalis | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 3. Streptococcus pneumoniae..... | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 4. Methicillin-resistant Staphylococcus Aureus (MRSA) | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 5. Staphylococcus aureus (not MRSA) | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 6. Mycoplasma pneumoniae | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 7. Pseudomonas Aureginosa | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 8. Chlamydomphila (or Chlamydia) pneumoniae | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 9. Oropharyngeal flora..... | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 10. Other_____ | | |

42. Was a blood culture done? No/NR 0 *Skip to 43* Yes 1

a. Culture Results Neg 0 *Skip to 43* Pos 1 Not Recorded 9 *Skip to 43*

- b. If yes, were any of the following reported in the blood culture?
- | | <u>No</u> | <u>Yes</u> |
|---|----------------------------|----------------------------|
| 1. Haemophilus Influenzae..... | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 2. Moraxella Catarrhalis | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 3. Streptococcus pneumoniae..... | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 4. Methicillin-resistant Staphylococcus Aureus (MRSA) | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 5. Staphylococcus aureus (not MRSA) | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 6. Other_____ | | |

43. Influenza swab Neg 0 Pos 1 Not Recorded 9

F. TREATMENTS / MEDICATIONS

- | | <u>No/NR</u> | <u>Yes</u> |
|---|----------------------------|----------------------------|
| 44. CPAP or BiPap | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 45. Mechanical Ventilation | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 46. Inhaled short-acting beta-agonists (ie,albuterol, xopenex) | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 47. Inhaled short-acting anticholinergics (ie, atrovent, ipratropium) | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 48. Nebulized Bronchodilators | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 49. Magnesium injections in ED..... | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 50. Oxygen (continuous or prn) | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 51. IV Antibiotics | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 52. Systemic Corticosteroid (IV or PO)..... | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| 53. IV Lasix or Furosemide | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |

	<u>At onset of time of event</u>		<u>At discharge</u>	
	No/NR	Yes	No/NR	Yes
54. Antibiotics-oral	a. 0 <input type="checkbox"/>	<input type="checkbox"/> 1 b. 0 <input type="checkbox"/>	1 <input type="checkbox"/>
55. Systemic corticosteroid (ie prednisone)	a. 0 <input type="checkbox"/>	<input type="checkbox"/> 1 b. 0 <input type="checkbox"/>	1 <input type="checkbox"/>
56. Inhaled short acting beta-agonists (ie albuterol)	a. 0 <input type="checkbox"/>	<input type="checkbox"/> 1 b. 0 <input type="checkbox"/>	1 <input type="checkbox"/>
57. Inhaled long-acting beta-agonist (ie, serevent)	a. 0 <input type="checkbox"/>	<input type="checkbox"/> 1 b. 0 <input type="checkbox"/>	1 <input type="checkbox"/>
58. Inhaled short-acting anticholinergics (ie, atrovent)	a. 0 <input type="checkbox"/>	<input type="checkbox"/> 1 b. 0 <input type="checkbox"/>	1 <input type="checkbox"/>
59. Inhaled long-acting anticholinergics	a. 0 <input type="checkbox"/>	<input type="checkbox"/> 1 b. 0 <input type="checkbox"/>	1 <input type="checkbox"/>
60. Inhaled corticosteroids	a. 0 <input type="checkbox"/>	<input type="checkbox"/> 1 b. 0 <input type="checkbox"/>	1 <input type="checkbox"/>
61. Nebulized bronchodilators	a. 0 <input type="checkbox"/>	<input type="checkbox"/> 1 b. 0 <input type="checkbox"/>	1 <input type="checkbox"/>
62. Leukotriene antagonist.....	a. 0 <input type="checkbox"/>	<input type="checkbox"/> 1 b. 0 <input type="checkbox"/>	1 <input type="checkbox"/>
63. Home oxygen	a. 0 <input type="checkbox"/>	<input type="checkbox"/> 1 b. 0 <input type="checkbox"/>	1 <input type="checkbox"/>

APPENDIX E Reviewer Forms

Myocardial Infarction diagnosis form (MID)

Heart Failure diagnosis form (HFD)

Pulmonary diagnosis form (PLD)



HCHS/SOL HEART FAILURE (HFD) DIAGNOSIS FORM

ID NUMBER:

FORM CODE: HFD
VERSION: A
8/17/2011

Contact Occasion SEQ #

ADMINISTRATIVE INFORMATION

0a. Completion Date: / /
Month Day Year

0b. Reviewer ID:

0c. Event ID:

0d. Event Date: / /

- | | No | Yes | Unknown |
|---|----------------------------|----------------------------|----------------------------|
| 1. Does this patient have a history of heart failure | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> | 9 <input type="checkbox"/> |
| 2. HF diagnosed by provider <u>AND</u> treatment provided for HF? | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> | 9 <input type="checkbox"/> |
| 3. Pulmonary edema/congestion on Chest X-ray | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> | 9 <input type="checkbox"/> |

- | Historical or imaging evidence of: | No | Yes, history | Yes, current imaging | Unknown |
|--|----------------------------------|----------------------------------|----------------------------------|------------------------------------|
| 4. Dilated ventricle? | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> | 2 <input type="checkbox"/> | 9 <input type="checkbox"/> |
| 5. Poor LV function (e.g., low EF or wall motion abnormalities) | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> | 2 <input type="checkbox"/> | 9 <input type="checkbox"/> |
| 6. Poor RV function | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> | 2 <input type="checkbox"/> | 9 <input type="checkbox"/> |
| 7. Diastolic dysfunction | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> | 2 <input type="checkbox"/> | 9 <input type="checkbox"/> |
| 8. Quantitative EF during this hospitalization (or within 3 months)? | | | | |
| 1 <input type="checkbox"/> ≥ 50 | 2 <input type="checkbox"/> 40-49 | 3 <input type="checkbox"/> 30-30 | 4 <input type="checkbox"/> 20-29 | 5 <input type="checkbox"/> <20 |
| | | | | 6 <input type="checkbox"/> unknown |

REVIEWER CLASSIFICATION

9. Does this patient have acute decompensated heart failure (ADHF)?
1 Definite 2 Probable 3 No, *skip to 10* 4 Unknown, *skip to 10*

If Definite or Probable ADHF then answer 9a-9c:

9a. How would you classify the severity of the exacerbation?

- 1 Mild 2 Moderate 3 Severe 4 unknown

9b. Was ADHF predominantly right-sided HF (normal LV EF)? 0 No 1 Yes 9 Unknown

9c. Were any of the following problems co-morbid with this event and could have precipitated this event?

- | | NO/NR | YES |
|---|--|--|
| c.1. Myocardial infarction | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| c.2. Atrial fibrillation/atrial flutter | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| c.3. Other arrhythmia | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| c.4. Fluid or volume overload | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| c.5. Medication noncompliance | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| c.6. Pulmonary embolus | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| c.7. Renal Insufficiency or failure | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| c.8. Cardiovascular procedure/surgery | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| c.9. Non-cardiovascular procedure/surgery | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| c.10. Pulmonary disease | 0 <input type="checkbox"/> | 1 <input type="checkbox"/> |
| c.11. Uncontrolled Hypertension | 0 <input type="checkbox"/> <i>skip to 11</i> | 1 <input type="checkbox"/> <i>skip to 11</i> |

10. If **not** definite or probable ADHF, then does this patient have Asymptomatic LV dysfunction (EF < 50%)?
0 No/NR 1 Yes 9 Unknown

11. Reviewer comments: _____



HCHS/SOL PULMONARY DIAGNOSIS (PLD) FORM

ID NUMBER:

FORM CODE: PLD
VERSION: A 7/20/11

Contact Occasion SEQ #

Administrative Information

0A. Completion Date: / /
Month Day Year

0B. Reviewer ID:

0C. Event ID:

0D. Event Date: / /
Month Day Year

	Yes		No		Unknown
	Definite	Probable	Probably Not	Definitely not	
1. Does this patient meet SOL criteria for a history of chronic lower respiratory disease (CLRD)?	1 <input type="checkbox"/>	2 <input type="checkbox"/>	3 <input type="checkbox"/> Go to Q2	4 <input type="checkbox"/> Go to Q2	5 <input type="checkbox"/> Go to Q2

If YES (definite or probable.)
Does this patient have a history of any of the following?

a. COPD.....	1 <input type="checkbox"/>	2 <input type="checkbox"/>	3 <input type="checkbox"/>	4 <input type="checkbox"/>	5 <input type="checkbox"/>
b. Emphysema	1 <input type="checkbox"/>	2 <input type="checkbox"/>	3 <input type="checkbox"/>	4 <input type="checkbox"/>	5 <input type="checkbox"/>
c. Chronic Bronchitis.....	1 <input type="checkbox"/>	2 <input type="checkbox"/>	3 <input type="checkbox"/>	4 <input type="checkbox"/>	5 <input type="checkbox"/>
d. Asthma.....	1 <input type="checkbox"/>	2 <input type="checkbox"/>	3 <input type="checkbox"/>	4 <input type="checkbox"/>	5 <input type="checkbox"/>

2. Is there evidence of other lung disease? No=0 Yes=1

a. If yes, specify lung disease _____

	Yes		No		Unknown
	Definite	Probable	Probably Not	Definitely not	
3. Does this patient have an exacerbation of underlying chronic lower respiratory disease (CLRD)?.....	1 <input type="checkbox"/>	2 <input type="checkbox"/>	3 <input type="checkbox"/>	4 <input type="checkbox"/>	5 <input type="checkbox"/>

a. If Yes (definite or probable), then which type of CLRD is the cause of this exacerbation? (select one)

1. Asthma predominant 3. Either asthma or COPD
2. COPD predominant 4. Unclassifiable

	Yes		No		Unknown/ unclassified
	Definite	Probable	Probably Not	Definitely not	
4. Did this patient have pneumonia (new infiltrate on chest imaging)?	1 <input type="checkbox"/>	2 <input type="checkbox"/>	3 <input type="checkbox"/>	4 <input type="checkbox"/>	5 <input type="checkbox"/>

5. Comments: _____

Appendix F - Event Summary Forms

Myocardial Infarction Event Summary
Heart Failure Event Summary
Pulmonary Event Summary

Event Summary Form (ESF) for Myocardial Infarction

EVENT ID: X050000600101

HCHS ID:	Gender:	Age at Baseline:	Age at Event:
X0000060	Female	54	55
Prevalent CHD without Angina?		No	
Prevalent CHD with Angina?		Yes	

Abnormal ECG:

Prevalent MI by ECG?	No
Prevalent Q wave?	No

Self-report hx of:

Angina [MHEA3]	Yes
MI [MHEA4]	No
Coronary bypass, angio, or stent [MHEA9]	No
CHF [MHEA5]	No
Stroke [MHEA10]	No
Carotid endarterectomy, angioplasty, or surgery [MHEA12]	No
PAD [MHEA14]	No
Leg angio, stent, or amputation for vasc [MHEA15]	No

Event Date Date of arrival: [mm/dd/yyyy] Date of discharge: [mm/dd/yyyy]

ICD-9 Discharge Codes:

Was the patient alive at discharge?

Signs and symptoms

Acute episode(s) of pain or discomfort (eg: tightness) in the chest, arm, shoulder, throat or jaw, either within 72 hours?

Did this pain or discomfort specifically involve the chest?

Was the pain or discomfort diagnosed as having a non-cardiac origin?

Doctor's diagnosis

Did a physician indicate any of these as being present during the hospitalization?

- Angina
- Acute myocardial Infarction
- Congestive heart failure exacerbation or pulmonary edema
- Shock or cardiogenic shock
- Ventricular fibrillation, cardiac arrest or asystole
- ST elevation > 1mm with pain that is not present on ECG without pain

History

Prior to this event was there history of any of the following:

- Myocardial infarction
 - MI within 4 weeks?
- Angina
- Percutaneous coronary intervention (PCI)
- CABG
- Coronary artery disease (CAD)
- Heart failure

Treatment/Tests

Coronary reperfusion (CABG, PCI, thrombolysis) attempted within 24 hours?

Any of the following treatments given during this hospitalization?

Coronary artery bypass graft surgery (CABG)

Coronary atherectomy

Intra-arterial or intravenous thrombolytic

Coronary angioplasty without stent

Coronary angioplasty with stent placement

Valve surgery

Non-cardiac surgery

Aortic balloon pump

Pacemaker placement (temporary or permanent)

Cardioversion or defibrillation

Were any 12 lead ECGs taken during this admission?

First ECG Date: Copy enclosed?

Second ECG Date: Copy enclosed?

Last ECG Date: Copy enclosed?

Third ECG Date: Copy enclosed?

Biomarkers

Mention of trauma, procedure, or rhabdomyolysis, one week prior to biomarkers?

Evidence of hemolytic disease during the hospitalization?

Active liver disease?

Biomarker set	Date	Total CK	CK-MB	Troponin	Type of Troponin
1 st set					
2 nd set					
3 rd set					
4 th set					
5 th set					
6 th set					
Lab Standards					
Upper Limits					
Normal					
Units					

If quantitative biomarker value not available and only qualitative value then answers "A-C" are:

A = negative/absent/normal,

B = weak positive/weak present/trace/ high-normal/small,

C = present/ positive/abnormal/medium/large

Event Summary Form (ESF) for Heart Failure

EVENT ID: X00000080101

HCHS ID:	Gender:	Age at Baseline:	Age at Event:
X0000008	Male	57	58

Self-report hx of:

CHF [MHEA5]	No
Angina [MHEA3]	No
MI [MHEA4]	No
Coronary bypass, angioplasty, or stent [MHEA9]	No
Prevalent CHD without Angina?	Yes
Prevalent CHD with Angina?	Yes

Event Date Date of arrival: [mm/dd/yyyy] Date of discharge: [mm/dd/yyyy]

ICD-9 Discharge Codes:

Was the patient alive at discharge?

Signs and symptoms

- An increase or new onset of paroxysmal nocturnal dyspnea (PND)
- An increase or new onset of orthopnea
- An increase or new onset of shortness of breath
- An increase or new onset edema
- Hypoxia
- Dyspnea (at rest)
- Dyspnea (walking or on exertion)
- New onset or progressive signs/symptoms of HF prior to presentation?

History

Prior to this event was there a history of any of the following:

- Diagnosis of heart failure
- Prior hospitalization for heart failure
- Treatment for heart failure

Physician Diagnosis

Evidence that event was exacerbation of HF?

Ejection Fraction

Prior to this hospitalization:
During this hospitalization:
TTE
TEE
Coronary angiography
Radionuclide ventriculogram (RVG)

Chest X-ray

Did the patient have any of the following signs on chest x-ray at any time during this event?

- Pulmonary edema or pulmonary vascular congestion
- Cardiomegaly or Cardiothoracic ratio ≥ 0.5
- Bilateral pleural effusion
- Unilateral pleural effusion

BNP Levels (include worst, upper limit normal)

BNP (pg/mL)

ProBNP (pg/mL)

Treatment with IV diuretic

Event Summary Form (ESF) for Pulmonary Diagnosis

EVENT ID: X000000100101

HCHS ID:	Gender:	Age at Baseline:	Age at Event:	Height	Predicted PEF	BMI
X0000010	F	55	556	161.00	27.9	

Event Date 12/28/2009 **Date of arrival:** [mm/dd/yyyy] **Date of discharge:** [mm/dd/yyyy]

ICD-9 Discharge Codes:

786.50 486 493.20 99.21

ER, hospital or observation:

Doctor’s diagnosis

Reason in doctor’s notes that this event may be exacerbation of COPD:

Reason in doctor’s notes that event may be exacerbation of asthma:

Signs and symptoms

Any of the following signs or symptoms at the time of the event:

- New onset or increase in cough:
- New onset or increase in sputum production:
- New onset or increase in sputum purulence:
- New onset or increase in wheezing:
- New onset or increase in dyspnea:
- Crackles / rales:
- Wheezing or rhonchi:

History

History of any of the following:

- Asthma:
- Emphysema:
- Chronic obstructive pulmonary disease(COPD):

Tests

Any of the following signs on chest x-ray at any time during this event?

- Hyperinflation:
- Flattened diaphragms:
- Emphysema:
- Infiltrate/consolidation:
- Pulmonary edema:
- Pleural effusion:

Prior PFTS: Percent predicted FEV₁: FEV₁/FVC:

Peak flows

Date of first PEF(R) taken after hospital/ED arrival: (mm/dd/yyyy)

First PEF recording (L/min)

Worst PEF recording (L/min) during this hospitalization

TREATMENTS / MEDICATIONS

- Inhaled short-acting beta-agonists (ie,albuterol, xopenex):
- Inhaled short-acting anticholinergics (ie, atrovent, ipratropium):
- Systemic corticosteroid (oral or IV):
- CPAP or BiPap:
- Mechanical Ventilation:

No/NR 0 Yes 1